

Appendix M-3 POWER Computer Model Results



PROJECTION OF WILDLIFE ECOLOGICAL RESPONSE (POWER)

NELSON, POPE & VOORHIS, LLC, MICROCOMPUTER MODEL

SPECIES LIST

INTRODUCTION

This appendix has been included to present the results of a computer model used to investigate the various wildlife species which can be expected to be found on the site considering the habitats established. This model was developed by and for the use of Nelson, Pope & Voorhis, LLC using available information and references for the various species. The model utilizes Excel spreadsheets to identify wildlife species commonly found in various Long Island habitats, based upon thorough research of available literature. The habitats investigated consisted of Pine Barrens, Successional Woodland, Successional Shrubland, Successional Old Field, Wooded Swamp, and Terrestrial Cultural (lawn/landscaping). Some of the species listed in this model would not be expected on the property given the surrounding development, but are present in similar habitats.

The first column identifies the common name of the species, presented with the main common name in alphabetical order (for example: red-tailed hawk would come before blue jay). The scientific name of particular species is in the second column. The third column shows the legal status of the species, of which there are four possible entries (Endangered, Threatened, Special Concern and Local Concern). The fourth column indicates the seasons during which the species might be expected to be present and the fifth column, of particular importance to the environmental setting, contains information on frequency of the species in the habitat (abundant, common, rare and non expected); the species activity in the habitat (nesting, hunting and resting). References are provided with the reference list provided at the end of the appendix. The printout contained in this appendix, coupled with the discussions provided in the main body of the report, provides significant information of the wildlife found, or expected to be found on site.

Pine Barren Species - Inventory and Characterisitics

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	Reference
			winter	spring	summer	fall		
common bobwhite	<i>Colinus virginianus</i>	none	X	X	X	X	R / N,F	4 8
gray catbird	<i>Dumetella carolinensis</i>	none		Late	X		R / N,F	4 9
black capped chickadee	<i>Parus atricapillus</i>	none	X	X	X	X	A / N,F / R	4 11
American crow	<i>Corvus brachyrhynchos</i>	none	X	X	X	X	A / N,H	4 11
house finch	<i>Carpodacus mexicanus</i>	none		X	X	X	A / N,F	4 20
purple finch	<i>Carpodacus purpureus</i>	none	X	X	X	X	C / N,F	4 20
common flicker	<i>Colaptes auratus</i>	none	X	X	X	X	C / N,F	4 14
common grackle	<i>Quiscalus quiscula</i>	none	X	X	X	X	C / N,F	4 6
ruffed grouse	<i>Bonasa umbellus</i>	none	X	X	X	X	R / N,F	4 8
broad-winged hawk	<i>Buteo platypterus</i>	none		X	X		R / N,H	4 16
Cooper's hawk	<i>Accipiter cooperii</i>	special concern		X	X		N / N,H	4 17
red-tailed hawk	<i>Buteo jamaicensis</i>	none	X	X	X	X	C / N,H	4 16
sharp-shinned hawk	<i>Accipiter striatus</i>	special concern	X	X	X	X	R / N,H	4 16
blue jay	<i>Cyanocitta cristata</i>	none	X	X	X	X	A / N,F	4 10
Northern (dark-eyed) junco	<i>Junco hyemalis</i>	none		X	X	Early	R / N,F	4 21
Eastern kingbird	<i>Tyrannus tyrannus</i>	none		X	X	Early	C / N,F	4 15
golden-crowned kinglet	<i>Regulus satrapa</i>	none	X	X		X	R / N,H	4 7
ruby-crowned kinglet	<i>Regulus calendula</i>	none	X	X		X	R / N,H	4 7
white-breasted nuthatch	<i>Sitta carolinensis</i>	none	X	X	X	X	A / N,F	4 9
ovenbird	<i>Seiurus aurocapillus</i>	none		Late	X	Early	C / N,F	4 19
great-horned owl	<i>Bubo virginianus</i>	none	X	X	X	X	C / N,H	4 17
long-eared owl	<i>Asio otus</i>	none	X	X	X	X	C / N,H	4 17
yellow-bellied sapsucker	<i>Sphyrapicus varius</i>	none		Late	X	Early	R / N,F	14
pine siskin	<i>Carduelis pinus</i>	none	X	X	X	X	N / N,F	4 20
chipping sparrow	<i>Spizella passerina</i>	none	X	X	X	X	R / N,F	4 21
fox sparrow	<i>Passerella iliaca</i>	none	X	X		X	R / F	20 21
song sparrow	<i>Melospiza melodia</i>	none	X	X	X	X	R / N,F	4 22
European starling	<i>Sturnus vulgaris</i>	none	X	X	X	X	C / N,F	4 23
chimney swift	<i>Chaetura pelagica</i>	none		X	X		C / F	4 42
scarlet tanager	<i>Piranga olivacea</i>	none		X	X		C / N,F	4
brown thrasher	<i>Toxostoma rufum</i>	none		X	X	Early	R / N,F	4 9

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	Reference
			winter	spring	summer	fall		
hermit thrush	<i>Catharus guttatus</i>	none	X	X	X	X	C / N,F	4 7
wood thrush	<i>Hylocichla mustelina</i>	none		X	X	Early	R / N,F	4 7
pine warbler	<i>Dendroica pinus</i>	none		X	X	Early	C / N,F	4 19
prairie warbler	<i>Dendroica discolor</i>	none		Late	X	Early	C / N,F	4 19
yellow-rumped warbler	<i>Dendroica coronata</i>	none		X	X	Early	C / N,F	4 8
cedar waxwing	<i>Bombycilla cedrorum</i>	none		X	X	Early	R / N,F	4 23 32
whip-poor-will	<i>Caprimulgus vociferous</i>	special concern		Late	X		C / N	4 12
Eastern wood-peewee	<i>Contopus virens</i>	none		X	X		C / N,F	4 15
red-bellied woodpecker	<i>Melanerpes carolinus</i>	none	X	X	X	X	R / N,F	4 14
house wren	<i>Troglodytes aedon</i>	none		Late	X	Early	R / N,F	4 9
Mammals								
big-brown bat	<i>Eptesicus fuscus</i>	none	X	X	X	X	R / N,F	1 29
hoary bat	<i>Lasiurus borealis</i>	none			Late	Early	C / N,F	45
Keen's bat	<i>Myotis keenii</i>	none			X	Early	R / N	1 29
little-brown bat	<i>Myotis lucifugus</i>	none	X	X	X	X	C / N,F	1 29
red bat	<i>Lasiurus borealis</i>	none		Late	X	Early	C / N,F	1 29
Eastern pipistrelle	<i>Pipistrellus subflavus</i>	none		X	X	Early	R / N,F	1 29
silver-haired bat	<i>Lasionycteris noctivagans</i>	none			X		R / N,F	1 29
Eastern chipmunk	<i>Tamias striatus</i>	none	X	X	X	X	R / N,F	1 29
Eastern cottontail	<i>Sylvilagus floridanus</i>	none	X	X	X	X	C / N,F	1 29
white-tailed deer	<i>Odocoileus virginianus</i>	none	X	X	X	X	C / N,F	1 25 29
red fox	<i>Vulpes vulpes</i>	none	X	X	X	X	C / N,H	1 29
Eastern mole	<i>Scalopus aquaticus</i>	none	X	X	X	X	C / N,F	1 29
house mouse	<i>Mus musculus</i>	none	X	X	X	X	N / N,F	1 29
meadow-jumping mouse	<i>Zapus hudsonicus</i>	none	X	X	X	X	R / N,F	1 29
white-footed mouse	<i>Peromyscus leucopus</i>	none	X	X	X	X	C / N,F	1 29
Virginia opossum	<i>Didelphis virginiana</i>	none	X	X	X	X	C / N,F	1 29
raccoon	<i>Procyon lotor</i>	none	X	X	X	X	C / N,F	1 29
masked shrew	<i>Sorex cinereus</i>	none	X	X	X	X	C / N,F	1 29
short-tailed shrew	<i>Blarina breuicauda</i>	none	X	X	X	X	A / N,F	1 29
striped skunk	<i>Mephitis mephitis</i>	none	X	X	X	X	N / N,F	1 29
Eastern gray squirrel	<i>Sciurus carolinensis</i>	none	X	X	X	X	C / N,F	1 29
southern-flying squirrel	<i>Glaucimys volans</i>	none	X	X	X	X	C / N,F	1 29
pine vole	<i>Microtus pinetorum</i>	none	X	X	X	X	C / N,F	1 29

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	Reference
			winter	spring	summer	fall		
long-tailed weasel	<i>Mustela frenata</i>	none	X	X	X	X	R / N,H	1 29
woodchuck	<i>Marmota monax</i>	none	X	X	X	X	R / N,F	1 29
Herptiles								
common gray treefrog	<i>Hyla versicolor</i>	none	X	X	X	X	C / N,F	33 37
red-backed salamander	<i>Plethodon cinereus cinereus</i>	none	X	X	X	X	R / N,F	34 36
Eastern tiger salamander	<i>Ambystoma tigrinum tigrinum</i>	endangered	X	X	X	X	R / F	36 38
marbled salamander	<i>Ambystoma opacum</i>	special concern	X	X	X	X	R / N,F	34 36 38
Eastern garter snake	<i>Thamnophis sirtalis</i>	none	X	X	X	X	C / N,F	38 40
eastern hognose snake	<i>Heterodon platyrhinos</i>	special concern	X	X	X	X	R / N,H	38
eastern milk snake	<i>Lampropeltis d. triangulum</i>	none	X	X	X	X	C / N,F	38 39
Eastern spadefoot toad	<i>Scaphiopus holbrooki</i>	special concern	X	X	X	X	C / N,F	33
Fowler's toad	<i>Bufo woodhousei fowleri</i>	none	X	X	X	X	C / F	33 37
Eastern box turtle	<i>Terrepepe carolina</i>	special concern	X	X	X	X	C / N,F	41

KEY:

Frequency:

A- abundant

C- common

R- rare

N- not expected

Activity:

N- nesting

H- hunting

R- resting

F- foraging

Successional Woodland Species - Inventory and Characteristics

Common Name	Scientific Name	Status	Found During				Frequency / Habitat Use	References
			winter	spring	summer	fall		
Birds	Birds							
gray catbird	<i>Dumetella carolinensis</i>	none		Late	X		C / N,F	4 9
black capped chickadee	<i>Parus atricapillus</i>	none	X	X	X	X	A / N,F	4 11
brown-headed cowbird	<i>Molothrus ater</i>	none		X	X	Early	A / N,F	4 6
brown creeper	<i>Certhia familiaris</i>	none		X	X	Early	C / N,F	4 9
American crow	<i>Corvus brachyrhynchos</i>	none	X	X	X	X	A / N,H	4 11
yellow-billed cuckoo	<i>Coccyzus americanus</i>	none		Late	X		C / N,F	4 12
mourning dove	<i>Zenaida macroura</i>	none	X	X	X	X	C / N,H	4 8
rock dove	<i>Columba livia</i>	none	X	X	X	X	C / N,F	4 8
house finch	<i>Carpodacus mexicanus</i>	none		X	X	X	A / N,F	4 20
common flicker	<i>Colaptes auratus</i>	none	X	X	X	X	A / N,F	4 14
common grackle	<i>Quiscalus quiscula</i>	none	X	X	X	X	A / N,F	4 6
ruffed grouse	<i>Bonasa umbellus</i>	none	X	X	X	X	R / N,F	4 8
rose-breasted grosbeak	<i>Pheucticus ludovicianus</i>	none		Late	X	Early	R / N,F	4 20
Cooper's hawk	<i>Accipiter cooperii</i>	special concern		X	X		N / N,H	4 17
red-tailed hawk	<i>Buteo jamaicensis</i>	none	X	X	X	X	C / H	4 16
sharp-shinned hawk	<i>Accipiter striatus</i>	special concern	X	X	X	X	N / N,F	4 16
blue jay	<i>Cyanocitta cristata</i>	none	X	X	X	X	A / N,F	4 10
Northern (dark-eyed) junco	<i>Junco hyemalis</i>	none	X			Late	C / N,F	4 21
American kestrel	<i>Falco sparverius</i>	none	X	X	X	X	C / N,H	4 17
Eastern kingbird	<i>Tyrannus tyrannus</i>	none		X	X	Early	C / N,F	4 15
golden-crowned kinglet	<i>Regulus satrapa</i>	none	X	X		X	R / N,H	4 7
ruby-crowned kinglet	<i>Regulus calendula</i>	none	X	X		X	R / N,H	4 7
Northern mockingbird	<i>Mimus polyglottos</i>	none	X	X	X	X	C / N,F	4 9
great-horned owl	<i>Bubo virginianus</i>	none	X	X	X	X	C / N,H	4 17
long-eared owl	<i>Asio otus</i>	none	X	X	X	X	C / N,H	4 17
American redstart	<i>Setophaga ruticilla</i>	none		Late	X	X	C / N,F	4 19
American robin	<i>Turdus migratorius</i>	none		X	X	Early	A / N,F	4 7
yellow-bellied sapsucker	<i>Sphyrapicus varius</i>	none		Late	X	Early	C / N,F	14
fox sparrow	<i>Passerella iliaca</i>	none	X	X		X	R / F	20 21
house sparrow	<i>Passer domesticus</i>	none	X	X	X	X	C / N,F	4 20
song sparrow	<i>Melospiza melodia</i>	none	X	X	X	X	A / N,F	4 22

Common Name	Scientific Name	Status	Found During				Frequency / Habitat Use	References
			winter	spring	summer	fall		
white-throated sparrow	<i>Zonotrichia albicollis</i>	none	X	X	X	X	C / N,F	4 22
European starling	<i>Sturnus vulgaris</i>	none	X	X	X	X	A / N,F	4 23
barn swallow	<i>Hirundo rustica</i>	none		Late	X		C / N,F	4 15
brown thrasher	<i>Toxostoma rufum</i>	none		X	X	Early	C / N,F	4 9
hermit thrush	<i>Catharus guttatus</i>	none	X	X	X	X	R / N,F	4 7
wood thrush	<i>Hylocichla mustelina</i>	none		X	X	Early	C / N,F	4 7
rufous-sided towhee	<i>Pipilo erythrophthalmus</i>	none		Late	X	Early	A / N,F	4 20
red-eyed vireo	<i>Vireo olivaceus</i>	none		Late	X		C / N,F	4 23
black-and-white warbler	<i>Mniotilta varia</i>	none		X	X		C / N,F	4 18
blue-winged warbler	<i>Vermivora pinus</i>	none		Late	X		C / N,F	4 14
chestnut-sided warbler	<i>Dendroica pensylvanica</i>	none		Late	X		C / N,F	4 19
cedar waxwing	<i>Bombycilla cedrorum</i>	none		X	X	Early	C / N,F	4 23 32
whip-poor-will	<i>Caprimulgus vociferous</i>	special concern		Late	X		C / N	4 12
Eastern wood-peewee	<i>Contopus virens</i>	none		X	X		C / N,F	4 15
American woodcock	<i>Philhela minor</i>	none		X	X	X	R / N,F	4 30
downy woodpecker	<i>Picoides pubescens</i>	none	X	X	X	X	A / N,F	4 14
hairy woodpecker	<i>Picoides villosus</i>	none	X	X	X	X	R / N,F	4 14
red-bellied woodpecker	<i>Melanerpes carolinus</i>	none	X	X	X	X	R / N,F	4 14
Carolina wren	<i>Thryothorus ludovicianus</i>	none	X	X	X	X	C / N,F	4 9
house wren	<i>Troglodytes aedon</i>	none		Late	X	Early	C / N,F	4 9
big-brown bat	<i>Eptesicus fuscus</i>	none	X	X	X	X	C / N,F	1 29
hoary bat	<i>Lasiurus borealis</i>	none			Late	Early	C / N,F	45
Keen's bat	<i>Myotis keenii</i>	none			X	Early	R / N	1 29
Mammals								
little-brown bat	<i>Myotis lucifugus</i>	none	X	X	X	X	C / N,F	1 29
red bat	<i>Lasiurus borealis</i>	none		Late	X	Early	C / N,F	1 29
silver-haired bat	<i>Lasionycteris noctivagans</i>	none			X		R / N,F	1 29
Eastern chipmunk	<i>Tamias striatus</i>	none	X	X	X	X	C / N,F	1 29
Eastern cottontail	<i>Sylvilagus floridanus</i>	none	X	X	X	X	A / N,F	1 29
white-tailed deer	<i>Odocoileus virginianus</i>	none	X	X	X	X	C / F	1 25 29
red fox	<i>Vulpes vulpes</i>	none	X	X	X	X	C / N,H	1 29
Eastern mole	<i>Scalopus aquaticus</i>	none	X	X	X	X	C / N,F	1 29
meadow-jumping mouse	<i>Zapus hudsonicus</i>	none	X	X	X	X	R / N,F	1 29
white-footed mouse	<i>Peromyscus leucopus</i>	none	X	X	X	X	C / N,F	1 29

Common Name	Scientific Name	Status	Found During				Frequency / Habitat Use	References
			winter	spring	summer	fall		
Virginia opossum	<i>Didelphis virginiana</i>	none	X	X	X	X	C / N,F	1 29
raccoon	<i>Procyon lotor</i>	none	X	X	X	X	C / N,F	1 29
masked shrew	<i>Sorex cinereus</i>	none	X	X	X	X	C / N,F	1 29
short-tailed shrew	<i>Blarina brevicauda</i>	none	X	X	X	X	A / N,F	1 29
striped skunk	<i>Mephitis mephitis</i>	none	X	X	X	X	N / N,F	1 29
Eastern gray squirrel	<i>Sciurus carolinensis</i>	none	X	X	X	X	C / N,F	1 29
meadow vole	<i>Microtus pennsylvanicus</i>	none	X	X	X	X	R / N,F	29 45
pine vole	<i>Microtus pinetorum</i>	none	X	X	X	X	C / N,F	1 29
long-tailed weasel	<i>Mustela frenata</i>	none	X	X	X	X	R / N,H	1 29
woodchuck	<i>Marmota monax</i>	none	X	X	X	X	R / N,F	1 29
Herptiles								
Eastern garter snake	<i>Thamnophis sirtalis</i>	none	X	X	X	X	C / N,F	38 40
eastern hognose snake	<i>Heterodon platyrhinos</i>	special concern	X	X	X	X	R / N,H	38
eastern milk snake	<i>Lampropeltis d. triangulum</i>	none	X	X	X	X	C / N,F	38 39

KEY:

Frequency:

- A- abundant
- C- common
- R- rare
- N- not expected

Activity:

- N- nesting
- H- hunting
- R- resting
- F- foraging

Successional Shrubland Species-Inventory and Characteristics

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	References
			Winter	Spring	Summer	Fall		
red-winged blackbird	<i>Agelaius phoeniceus</i>	none		X	X	Early	C / N,F	4 6
common bobwhite	<i>Colinus virginianus</i>	none	X	X	X	X	C / N,F	4 8
indigo bunting	<i>Passerina cyanea</i>	none		Late	X	Early	C / N,F	4 20
Northern cardinal	<i>Cardinalis cardinalis</i>	none	X	X	X	X	C / N,F	4 20
gray catbird	<i>Dumetella carolinensis</i>	none		Late	X		A / N,F	4 9
black capped chickadee	<i>Parus atricapillus</i>	none	X	X	X	X	A / N,F	4 11
brown-headed cowbird	<i>Molothrus ater</i>	none		X	X	Early	A / N,F	4 6
American crow	<i>Corvus brachyrhynchos</i>	none	X	X	X	X	A / H	4 11
black-billed cuckoo	<i>Coccyzus erythrophthalmus</i>	none		Late	X		C / N,F	4 11
yellow-billed cuckoo	<i>Coccyzus americanus</i>	none		Late	X		C / N,F	4 12
mourning dove	<i>Zenaida macroura</i>	none	X	X	X	X	A / N,H	4 8
American goldfinch	<i>Carduelis tristis</i>	none		X	X	X	C / N,F	4 20
common flicker	<i>Colaptes auratus</i>	none	X	X	X	X	R / H	4 14
least flycatcher	<i>Empidonax minimus</i>	none		Late	X		R / N,F	4 15
willow flycatcher	<i>Empidonax traillii</i>	none		Late	X		C / N,F	4 15
common grackle	<i>Quiscalus quiscula</i>	none	X	X	X	X	A / N,F	4 6
ruffed grouse	<i>Bonasa umbellus</i>	none	X	X	X	X	C / N,F	4 8
rose-breasted grosbeak	<i>Pheucticus ludovicianus</i>	none		Late	X	Early	C / N,F	4 20
Northern harrier	<i>Circus cyaneus</i>	threatened	X	X	X	X	R / H	4 16
Cooper's hawk	<i>Accipiter cooperii</i>	special concern		X	X		N / N,H	4 17
red-tailed hawk	<i>Buteo jamaicensis</i>	none	X	X	X	X	C / H	4 16
sharp-shinned hawk	<i>Accipiter striatus</i>	special concern	X	X	X	X	N / N,F	4 16
blue jay	<i>Cyanocitta cristata</i>	none	X	X	X	X	A / N,F	4 10
Northern (dark-eyed) junco	<i>Junco hyemalis</i>	none	X			Late	C / N,F	4 21
American kestrel	<i>Falco sparverius</i>	none	X	X	X	X	C / H	4 17
Eastern kingbird	<i>Tyrannus tyrannus</i>	none		X	X	Early	A / N,F	4 15
golden-crowned kinglet	<i>Regulus satrapa</i>	none	X	X		X	R / N,H	4 7
ruby-crowned kinglet	<i>Regulus calendula</i>	none	X	X		X	R / N,H	4 7
Eastern meadowlark	<i>Sturnella magna</i>	none		Late	X		C / N,F	4 6
Northern mockingbird	<i>Mimus polyglottos</i>	none	X	X	X	X	A / N,F	4 9
barn owl	<i>Tyto alba</i>	none	X	X	X	X	R / H	4 17
great-horned owl	<i>Bubo virginianus</i>	none	X	X	X	X	C / N,H	4 17

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	References
			winter	spring	summer	fall		
ring-necked pheasant	<i>Phasianus colchicus</i>	none	X	X	X	X	C / N,F	4 8
American redstart	<i>Setophaga ruticilla</i>	none		Late	X	X	C / N,F	4 19
American robin	<i>Turdus migratorius</i>	none		X	X	Early	A / N,F	4 7
pine siskin	<i>Carduelis pinus</i>	none	X	X	X	X	N / N,F	4 20
fox sparrow	<i>Passerella iliaca</i>	none	X	X		X	C / F	20 21
field sparrow	<i>Spizella pusilla</i>	none		X	X	X	C / N,F	4 21
grasshopper sparrow	<i>Ammodramus savannarum</i>	special concern		X	X	Early	C / N,F	4 20
song sparrow	<i>Melospiza melodia</i>	none	X	X	X	X	A / N,F	4 22
swamp sparrow	<i>Melospiza georgiana</i>	none	X	X	X	X	C / N,F	4 22
white-crowned sparrow	<i>Zonotrichia leucophrys</i>	none	X	X		X	C / F	22 32
white-throated sparrow	<i>Zonotrichia albicollis</i>	none	X	X	X	X	C / N,F	4 22
European starling	<i>Sturnus vulgaris</i>	none	X	X	X	X	A / N,F	4 23
barn swallow	<i>Hirundo rustica</i>	none		Late	X		C / N,F	4 15
brown thrasher	<i>Toxostoma rufum</i>	none		X	X	Early	C / N,F	4 9
hermit thrush	<i>Catharus guttatus</i>	none	X	X	X	X	R / N,F	4 7
wood thrush	<i>Hylocichla mustelina</i>	none		X	X	Early	C / N,F	4 7
rufous-sided towhee	<i>Pipilo erythrophthalmus</i>	none		Late	X	Early	A / N,F	4 20
white eyed vireo	<i>Vireo griseus</i>	none		Late	X		C / N,F	4 23
black-and-white warbler	<i>Mniotilta varia</i>	none		X	X		R / N,F	4 18
blue-winged warbler	<i>Vermivora pinus</i>	none		Late	X		C / N,F	4 14
chestnut-sided warbler	<i>Dendroica pensylvanica</i>	none		Late	X		C / N,F	4 19
prairie warbler	<i>Dendroica discolor</i>	none		Late	X	Early	C / N,F	4 19
cedar waxwing	<i>Bombycilla cedrorum</i>	none		X	X	Early	C / N,F	4 23 32
whip-poor-will	<i>Caprimulgus vociferous</i>	special concern		Late	X		C / F	4 12
American woodcock	<i>Philhela minor</i>	none		X	X	X	C / N,F	4 30
Carolina wren	<i>Thryothorus ludovicianus</i>	none	X	X	X	X	C / N,F	4 9
house wren	<i>Troglodytes aedon</i>	none		Late	X	Early	C / N,F	4 9
common yellowthroat	<i>Geothlypis trichas</i>	none		X	X	Early	C / N,F	4 19
Mammals								
Eastern chipmunk	<i>Tamias striatus</i>	none	X	X	X	X	C / N,F	1 29
Eastern cottontail	<i>Sylvilagus floridanus</i>	none	X	X	X	X	A / N,F	1 29
white-tailed deer	<i>Odocoileus virginianus</i>	none	X	X	X	X	C / F	1 25 29
red fox	<i>Vulpes vulpes</i>	none	X	X	X	X	C / N,H	1 29
Eastern mole	<i>Scalopus aquaticus</i>	none	X	X	X	X	C / N,F	1 29

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	References
			winter	spring	summer	fall		
house mouse	<i>Mus musculus</i>	none	X	X	X	X	R / N,F	1 29
meadow-jumping mouse	<i>Zapus hudsonicus</i>	none	X	X	X	X	R / N,F	1 29
white-footed mouse	<i>Peromyscus leucopus</i>	none	X	X	X	X	C / N,F	1 29
Virginia opossum	<i>Didelphis virginiana</i>	none	X	X	X	X	C / N,F	1 29
raccoon	<i>Procyon lotor</i>	none	X	X	X	X	C / F	1 29
black rat	<i>Rattus rattus</i>	none	X	X	X	X	R / N,F	1 29
short-tailed shrew	<i>Blarina breuicauda</i>	none	X	X	X	X	A / N,F	1 29
striped skunk	<i>Mephitis mephitis</i>	none	X	X	X	X	N / N,F	1 29
meadow vole	<i>Microtus pennsylvanicus</i>	none	X	X	X	X	C / N,F	29 45
pine vole	<i>Microtus pinetorum</i>	none	X	X	X	X	C / N,F	1 29
long-tailed weasel	<i>Mustela frenata</i>	none	X	X	X	X	R / N,H	1 29
woodchuck	<i>Marmota monax</i>	none	X	X	X	X	R / N,F	1 29
Herptiles								
Eastern garter snake	<i>Thamnophis sirtalis</i>	none	X	X	X	X	C / N,F	38 40
eastern hognose snake	<i>Heterodon platyrhinos</i>	special concern	X	X	X	X	R / N,H	38
eastern milk snake	<i>Lampropeltis d. triangulum</i>	none	X	X	X	X	C / N,F	38 39
Fowler's toad	<i>Bufo woodhousei fowleri</i>	none	X	X	X	X	C / F	33 37

KEY:

Frequency:

A- abundant
C- common
R- rare
N- not expected

Activity:

N- nesting
H- hunting
R- resting
F- foraging

Successional Old Field Species - Inventory and Characteristics

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	References
			Winter	Spring	Summer	Fall		
Birds								
red-winged blackbird	<i>Agelaius phoeniceus</i>	none		X	X	Early	C / N,F	4 6
Eastern bluebird	<i>Sialia sialis</i>	none		X	X	Early	R / N,F	4 7
common bobwhite	<i>Colinus virginianus</i>	none	X	X	X	X	C / N,F	4 8
indigo bunting	<i>Passerina cyanea</i>	none		Late	X	Early	C / N,F	4 20
Northern cardinal	<i>Cardinalis cardinalis</i>	none	X	X	X	X	C / N,F	4 20
gray catbird	<i>Dumetella carolinensis</i>	none		Late	X		A / N,F	4 9
black capped chickadee	<i>Parus atricapillus</i>	none	X	X	X	X	C / H	4 11
brown-headed cowbird	<i>Molothrus ater</i>	none		X	X	Early	A / H	4 6
American crow	<i>Corvus brachyrhynchos</i>	none	X	X	X	X	A / H	4 11
black-billed cuckoo	<i>Coccyzus erythrophthalmus</i>	none		Late	X		C / N,F	4 11
yellow-billed cuckoo	<i>Coccyzus americanus</i>	none		Late	X		C / N,F	4 12
mourning dove	<i>Zenaida macroura</i>	none	X	X	X	X	A / N,H	4 8
rock dove	<i>Columba livia</i>	none	X	X	X	X	A / N,F	4 8
American goldfinch	<i>Carduelis tristis</i>	none		X	X	X	C / N,F	4 20
house finch	<i>Carpodacus mexicanus</i>	none		X	X	X	A / N,F	4 20
common flicker	<i>Colaptes auratus</i>	none	X	X	X	X	C / N,F	4 14
least flycatcher	<i>Empidonax minimus</i>	none		Late	X		R / N,F	4 15
willow flycatcher	<i>Empidonax traillii</i>	none		Late	X		C / N,F	4 15
common grackle	<i>Quiscalus quiscula</i>	none	X	X	X	X	A / N,F	4 6
ruffed grouse	<i>Bonasa umbellus</i>	none	X	X	X	X	C / N,F	4 8
rose-breasted grosbeak	<i>Pheucticus ludovicianus</i>	none		Late	X	Early	C / N,F	4 20
Northern harrier	<i>Circus cyaneus</i>	threatened	X	X	X	X	R / H	4 16
Cooper's hawk	<i>Accipiter cooperii</i>	special concern		X	X		N / N,H	4 17
red-tailed hawk	<i>Buteo jamaicensis</i>	none	X	X	X	X	C / H	4 16
sharp-shinned hawk	<i>Accipiter striatus</i>	special concern	X	X	X	X	N / N,F	4 16
blue jay	<i>Cyanocitta cristata</i>	none	X	X	X	X	A / N,F	4 10
Northern (dark-eyed) junco	<i>Junco hyemalis</i>	none	X			Late	C / N,F	4 21
American kestrel	<i>Falco sparverius</i>	none	X	X	X	X	C / H	4 17
Eastern kingbird	<i>Tyrannus tyrannus</i>	none		X	X	Early	A / N,F	4 15
Eastern meadowlark	<i>Sturnella magna</i>	none		Late	X		C / N,F	4 6
Northern mockingbird	<i>Mimus polyglottos</i>	none	X	X	X	X	A / N,F	4 9

Common Name	Scientific Name	Status	Found During			Fall	Frequency/ Habitat Use	References
			Winter	Spring	Summer			
common nighthawk	<i>Chordeiles minor</i>	special concern		Late	X		R / N,F	4 12
barn owl	<i>Tyto alba</i>	none	X	X	X	X	R / H	4 17
great-horned owl	<i>Bubo virginianus</i>	none	X	X	X	X	C / N,H	4 17
ring-necked pheasant	<i>Phasianus colchicus</i>	none	X	X	X	X	C / N,F	4 8
American robin	<i>Turdus migratorius</i>	none		X	X	Early	A / N,F	4 7
chipping sparrow	<i>Spizella passerina</i>	none	X	X	X	X	C / N,F	4 21
fox sparrow	<i>Passerella iliaca</i>	none	X	X		X	C / F	20 21
field sparrow	<i>Spizella pusilla</i>	none		X	X	X	C / N,F	4 21
grasshopper sparrow	<i>Ammodramus savannarum</i>	special concern		X	X	Early	C / N,F	4 20
house sparrow	<i>Passer domesticus</i>	none	X	X	X	X	C / N,F	4 20
Savannah sparrow	<i>Passerculus sandwichensis</i>	none		X	X	Early	R / N,F	4 21
song sparrow	<i>Melospiza melodia</i>	none	X	X	X	X	A / N,F	4 22
swamp sparrow	<i>Melospiza georgiana</i>	none	X	X	X	X	C / N,F	4 22
white-crowned sparrow	<i>Zonotrichia leucophrys</i>	none	X	X		X	C / F	22 32
European starling	<i>Sturnus vulgaris</i>	none	X	X	X	X	A / N,F	4 23
barn swallow	<i>Hirundo rustica</i>	none		Late	X		C / N,F	4 15
brown thrasher	<i>Toxostoma rufum</i>	none		X	X	Early	C / N,F	4 9
rufous-sided towhee	<i>Pipilo erythrophthalmus</i>	none		Late	X	Early	A / N,F	4 20
black-and-white warbler	<i>Mniotilta varia</i>	none		X	X		R / N,F	4 18
blue-winged warbler	<i>Vermivora pinus</i>	none		Late	X		C / N,F	4 14
chestnut-sided warbler	<i>Dendroica pensylvanica</i>	none		Late	X		R / N,F	4 19
prairie warbler	<i>Dendroica discolor</i>	none		Late	X	Early	C / N,F	4 19
yellow warbler	<i>Dendrocica petchia</i>	none		Late	X		R / N,F	4 18
cedar waxwing	<i>Bombycilla cedrorum</i>	none		X	X	Early	C / N,F	4 23 32
whip-poor-will	<i>Caprimulgus vociferous</i>	special concern		Late	X		C / F	4 12
American woodcock	<i>Philhela minor</i>	none		X	X	X	C / N,F	4 30
red-headed woodpecker	<i>Melanerpes erythrocephalus</i>	special concern		X	X	Early	C / N,F	4 14
house wren	<i>Troglodytes aedon</i>	none		Late	X	Early	C / N,F	4 9
common yellowthroat	<i>Geothlypis trichas</i>	none		X	X	Early	C / N,F	4 19
Mammals								
Eastern chipmunk	<i>Tamias striatus</i>	none	X	X	X	X	C / N,F	1 29
Eastern cottontail	<i>Sylvilagus floridanus</i>	none	X	X	X	X	A / N,F	1 29
white-tailed deer	<i>Odocoileus virginianus</i>	none	X	X	X	X	C / F	1 25 29
red fox	<i>Vulpes vulpes</i>	none	X	X	X	X	C / H	1 29

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	References
			Winter	Spring	Summer	Fall		
Eastern mole	<i>Scalopus aquaticus</i>	none	X	X	X	X	C / N,F	1 29
house mouse	<i>Mus musculus</i>	none	X	X	X	X	R / N,F	1 29
meadow-jumping mouse	<i>Zapus hudsonicus</i>	none	X	X	X	X	R / N,F	1 29
white-footed mouse	<i>Peromyscus leucopus</i>	none	X	X	X	X	C / N,F	1 29
Virginia opossum	<i>Didelphis virginiana</i>	none	X	X	X	X	C / N,F	1 29
raccoon	<i>Procyon lotor</i>	none	X	X	X	X	C / F	1 29
black rat	<i>Rattus rattus</i>	none	X	X	X	X	R / N,F	1 29
Norway rat	<i>Rattus norvegicus</i>	none	X	X	X	X	C / N,F	1 29
least shrew	<i>Cryptotis parva</i>	none	X	X	X	X	N / N,F	1 29
short-tailed shrew	<i>Blarina breuicauda</i>	none	X	X	X	X	A / N,F	1 29
striped skunk	<i>Mephitis mephitis</i>	none	X	X	X	X	N / N,F	1 29
meadow vole	<i>Microtus pennsylvanicus</i>	none	X	X	X	X	C / N,F	29 45
pine vole	<i>Microtus pinetorum</i>	none	X	X	X	X	C / N,F	1 29
long-tailed weasel	<i>Mustela frenata</i>	none	X	X	X	X	R / N,H	1 29
woodchuck	<i>Marmota monax</i>	none	X	X	X	X	R / N,F	1 29
Herptiles								
Eastern garter snake	<i>Thamnophis sirtalis</i>	none	X	X	X	X	C / N,F	38 40
eastern hognose snake	<i>Heterodon platyrhinos</i>	special concern	X	X	X	X	R / N,H	38
eastern milk snake	<i>Lampropeltis d. triangulum</i>	none	X	X	X	X	C / N,F	38 39
Fowler's toad	<i>Bufo woodhousei fowleri</i>	none	X	X	X	X	C / F	33 37

KEY:

Frequency:

A- abundant

C- common

R- rare

N- not expected

Activity:

N- nesting

H- hunting

R- resting

F- foraging

Wooded Swamp Species: Inventory and Characteristics

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	Reference
			winter	spring	summer	fall		
Birds								
red-winged blackbird	<i>Agelaius phoeniceus</i>	none		X	X	Early	A / N,F	4 6
Northern cardinal	<i>Cardinalis cardinalis</i>	none	X	X	X	X	C / N,F	4 20
wood duck	<i>Aix sponsa</i>	none		X	X	Early	C / N,F	4 27
great-crested flycatcher	<i>Myiarchus crinitus</i>	none		Late	X		C / N,F	4 15
blue-grey gnatcatcher	<i>Poliophtila caerulea</i>	none		X	X		R / N,F	4 7
rose-breasted grosbeak	<i>Pheucticus ludovicianus</i>	none		Late	X	Early	R / N,F	4 20
yellow-crowned night-heron	<i>Nycticorax violaceus</i>	none	X	X	X	X	C / N,H	4 26
mallard	<i>Anas platyrhynchos</i>	none		X	X	X	C / N,F	4 27
white-breasted nuthatch	<i>Sitta carolinensis</i>	none	X	X	X	X	A / N,F	4 9
osprey	<i>Pandion haliaetus</i>	special concern		X	X	Early	C / N,H	4 16
great-horned owl	<i>Bubo virginianus</i>	none	X	X	X	X	C / N,H	4 17
saw-whet owl	<i>Aegolius acadicus</i>	none	X	X	X	X	C / N,H	4 17
Eastern phoebe	<i>Sayornis phoebe</i>	none		X	X	Early	C / N,F	4 15
spotted sandpiper	<i>Actitis macularia</i>	none		X	X	Early	R / N,F	4 31 32
swamp sparrow	<i>Melospiza georgiana</i>	none	X	X	X	X	C / N,F	4 22
European starling	<i>Sturnus vulgaris</i>	none	X	X	X	X	R / N,F	4 23
tree swallow	<i>Tachycineta bicolor</i>	none		X	X		C / N,F	4 15
tufted titmouse	<i>Parus bicolor</i>	none	X	X	X	X	C / N,F	4 11
veery	<i>Catharus fuscescens</i>	none		Late	X		C / N,F	4 7
yellow warbler	<i>Dendrocica petchia</i>	none		Late	X		C / N,F	4 18
American woodcock	<i>Philhela minor</i>	none		X	X	X	C / N,F	4 30
downy woodpecker	<i>Picooides pubescens</i>	none	X	X	X	X	A / N,F	4 14
hairy woodpecker	<i>Picooides villosus</i>	none	X	X	X	X	C / N,F	4 14
red-bellied woodpecker	<i>Melanerpes carolinus</i>	none	X	X	X	X	C / N,F	4 14
Mammals								
big-brown bat	<i>Eptesicus fuscus</i>	none	X	X	X	X	R / H	1 29
little-brown bat	<i>Myotis lucifugus</i>	none	X	X	X	X	R / H	1 29
Eastern cottontail	<i>Sylvilagus floridanus</i>	none	X	X	X	X	A / F	1 29
white-tailed deer	<i>Odocoileus virginianus</i>	none	X	X	X	X	C / F	1 25 29
red fox	<i>Vulpes vulpes</i>	none	X	X	X	X	C / H	1 29
mink	<i>Mustela vison</i>	none	X	X	X	X	C / N,F	1

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	Reference
			winter	spring	summer	fall		
star-nosed mole	<i>Condylura cristata</i>	none	X	X	X	X	R / N,F	1 29
meadow-jumping mouse	<i>Zapus hudsonicus</i>	none	X	X	X	X	R / N,F	1 29
white-footed mouse	<i>Peromyscus leucopus</i>	none	X	X	X	X	C / N,F	1 29
muskrat	<i>Ondarta zibethicus</i>	none	X	X	X	X	C / N,F	1 29
Virginia opossum	<i>Didelphis virginiana</i>	none	X	X	X	X	C / N,F	1 29
raccoon	<i>Procyon lotor</i>	none	X	X	X	X	C / N,F	1 29
Norway rat	<i>Rattus norvegicus</i>	none	X	X	X	X	C / N,F	1 29
masked shrew	<i>Sorex cinereus</i>	none	X	X	X	X	C / N,F	1 29
short-tailed shrew	<i>Blarina brevicauda</i>	none	X	X	X	X	A / N,F	1 29
Eastern gray squirrel	<i>Sciurus carolinensis</i>	none	X	X	X	X	A / N,F	1 29
southern-flying squirrel	<i>Glaucimys volans</i>	none	X	X	X	X	C / N,F	1 29
meadow vole	<i>Microtus pennsylvanicus</i>	none	X	X	X	X	R / N,F	29 45
pine vole	<i>Microtus pinetorum</i>	none	X	X	X	X	R / N,F	1 29
long-tailed weasel	<i>Mustela frenata</i>	none	X	X	X	X	R / N,H	1 29
Herptiles								
bull frog	<i>Rana catesbeiana</i>	none	X	X	X	X	C / N,H	33 34 35 37
common gray treefrog	<i>Hyla versicolor</i>	none	X	X	X	X	C / N,F	33 37
green frog	<i>Rana clamitans</i>	none	X	X	X	X	C / N,H	33 35 37
Southern leopard frog	<i>Rana pipiens sphenoccephala</i>	special concern	X	X	X	X	R / N,F	35 37 38
wood frog	<i>Rana sylvatica</i>	none	X	X	X	X	C / N,F	33 37
red-spotted newt	<i>Notophthalmus viridescens</i>	none	X	X	X	X	C / N,F	36 38
spring peeper	<i>Hyla crucifer</i>	none	X	X	X	X	A / N,F	33 35 38
spotted salamander	<i>Ambystoma maculatum</i>	none	X	X	X	X	C / N,F	34 36 38
Eastern tiger salamander	<i>Ambystoma tigrinum tigrinum</i>	endangered	X	X	X	X	R / N,F	36 38
marbled salamander	<i>Ambystoma opacum</i>	special concern	X	X	X	X	R / N,F	34 36 38
Eastern garter snake	<i>Thamnophis sirtalis</i>	none	X	X	X	X	C / N,F	38 40
eastern milk snake	<i>Lampropeltis d. triangulum</i>	none	X	X	X	X	C / N,F	38 39
Eastern ribbon snake	<i>Thamnophis s. sauritus</i>	none	X	X	X	X	C / N,F	38 40
Northern brown snake	<i>Storeria dekayi</i>	none	X	X	X	X	C / N,H	38
Northern ringneck snake	<i>Diadophis punctatus</i>	none	X	X	X	X	C / H	38
Northern water snake	<i>Natrix sipedon sipedon</i>	none	X	X	X	X	C / N,F	38 39
painted turtle	<i>Chrysemys picta</i>	none	X	X	X	X	N / N,F	38 40

KEY:

Frequency:

A- abundant
C- common
R- rare
N- not expected

Activity:

N- nesting
H- hunting
R- resting
F- foraging

Terrestrial Cultural Species - Inventory and Characteristics

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	References
			winter	spring	summer	fall		
Birds								
red-winged blackbird	<i>Agelaius phoeniceus</i>	none		X	X	Early	C / N,F	4 6
Eastern bluebird	<i>Sialia sialis</i>	none		X	X	Early	R / N,F	4 7
common bobwhite	<i>Colinus virginianus</i>	none	X	X	X	X	C / N,F	4 8
indigo bunting	<i>Passerina cyanea</i>	none		Late	X	Early	N / F	4 20
Northern cardinal	<i>Cardinalis cardinalis</i>	none	X	X	X	X	C / N,F	4 20
gray catbird	<i>Dumetella carolinensis</i>	none		Late	X		C / F	4 9
brown-headed cowbird	<i>Molothrus ater</i>	none		X	X	Early	A / H	4 6
American crow	<i>Corvus brachyrhynchos</i>	none	X	X	X	X	A / N,H	4 11
mourning dove	<i>Zenaidura macroura</i>	none	X	X	X	X	A / N,H	4 8
rock dove	<i>Columba livia</i>	none	X	X	X	X	A / N,F	4 8
great-crested flycatcher	<i>Myiarchus crinitus</i>	none		Late	X		C / N,F	4 15
common grackle	<i>Quiscalus quiscula</i>	none	X	X	X	X	A / N,F	4 6
Northern harrier	<i>Circus cyaneus</i>	threatened	X	X	X	X	R / H	4 16
Cooper's hawk	<i>Accipiter cooperii</i>	special concern		X	X		N / N,H	4 17
red-tailed hawk	<i>Buteo jamaicensis</i>	none	X	X	X	X	C / H	4 16
sharp-shinned hawk	<i>Accipiter striatus</i>	special concern	X	X	X	X	N / N,F	4 16
blue jay	<i>Cyanocitta cristata</i>	none	X	X	X	X	A / N,F	4 10
American kestrel	<i>Falco sparverius</i>	none	X	X	X	X	C / N,H	4 17
killdeer	<i>Charadrius vociferus</i>	none		X	X	X	C / N,F	4 31 32
Eastern kingbird	<i>Tyrannus tyrannus</i>	none		X	X	Early	A / N,F	4 15
Eastern meadowlark	<i>Sturnella magna</i>	none		Late	X		C / N,F	4 6
Northern mockingbird	<i>Mimus polyglottos</i>	none	X	X	X	X	A / N,F	4 9
common nighthawk	<i>Chordeiles minor</i>	special concern		Late	X		R / N,F	4 12
barn owl	<i>Tyto alba</i>	none	X	X	X	X	R / N,H	4 17
great-horned owl	<i>Bubo virginianus</i>	none	X	X	X	X	R / N,H	4 17
ring-necked pheasant	<i>Phasianus colchicus</i>	none	X	X	X	X	C / N,F	4 8
black-bellied plover	<i>Pluvialis squatarola</i>	none		Early	X	Early	R / F	31 32
American robin	<i>Turdus migratorius</i>	none		X	X	Early	A / N,F	4 7
chipping sparrow	<i>Spizella passerina</i>	none	X	X	X	X	C / N,F	4 21
field sparrow	<i>Spizella pusilla</i>	none		X	X	X	R / N,F	4 21
grasshopper sparrow	<i>Ammodramus savannarum</i>	special concern		X	X	Early	R / N,F	4 20

Common Name	Scientific Name	Status	Found During				Frequency/ Habitat Use	References
			winter	spring	summer	fall		
house sparrow	<i>Passer domesticus</i>	none	X	X	X	X	A / N,F	4 20
Savannah sparrow	<i>Passerculus sandwichensis</i>	none		X	X	Early	R / N,F	4 21
song sparrow	<i>Melospiza melodia</i>	none	X	X	X	X	C / N,F	4 22
white-crowned sparrow	<i>Zonotrichia leucophrys</i>	none	X	X		X	C / F	22 32
European starling	<i>Sturnus vulgaris</i>	none	X	X	X	X	A / N,F	4 23
barn swallow	<i>Hirundo rustica</i>	none		Late	X		A / N,F	4 15
chimney swift	<i>Chaetura pelagica</i>	none		X	X		C / F	4 42
brown thrasher	<i>Toxostoma rufum</i>	none		X	X	Early	R / N,F	4 9
chestnut-sided warbler	<i>Dendroica pensylvanica</i>	none		Late	X		N / N,F	4 19
cedar waxwing	<i>Bombycilla cedrorum</i>	none		X	X	Early	C / N,F	4 23 32
whip-poor-will	<i>Caprimulgus vociferous</i>	special concern		Late	X		C / F	4 12
American woodcock	<i>Philhela minor</i>	none		X	X	X	R / N,F	4 30
red-headed woodpecker	<i>Melanerpes erythrocephalus</i>	special concern		X	X	Early	C / N,F	4 14
house wren	<i>Troglodytes aedon</i>	none		Late	X	Early	C / N,F	4 9
Mammals								
Eastern pipistrelle	<i>Pipistrellus subflavus</i>	none		X	X	Early	N / F	1 29
Eastern cottontail	<i>Sylvilagus floridanus</i>	none	X	X	X	X	A / N,F	1 29
white-tailed deer	<i>Odocoileus virginianus</i>	none	X	X	X	X	C / F	1 25 29
red fox	<i>Vulpes vulpes</i>	none	X	X	X	X	C / H	1 29
house mouse	<i>Mus musculus</i>	none	X	X	X	X	C / N,F	1 29
meadow-jumping mouse	<i>Zapus hudsonicus</i>	none	X	X	X	X	R / N,F	1 29
white-footed mouse	<i>Peromyscus leucopus</i>	none	X	X	X	X	C / N,F	1 29
Virginia opossum	<i>Didelphis virginiana</i>	none	X	X	X	X	C / N,F	1 29
raccoon	<i>Procyon lotor</i>	none	X	X	X	X	C / F	1 29
black rat	<i>Rattus rattus</i>	none	X	X	X	X	N / N,F	1 29
Norway rat	<i>Rattus norvegicus</i>	none	X	X	X	X	C / N,F	1 29
least shrew	<i>Cryptotis parva</i>	none	X	X	X	X	N / N,F	1 29
striped skunk	<i>Mephitis mephitis</i>	none	X	X	X	X	N / N,F	1 29
pine vole	<i>Microtus pinetorum</i>	none	X	X	X	X	C / N,F	1 29
long-tailed weasel	<i>Mustela frenata</i>	none	X	X	X	X	R / N,H	1 29
Heptiles								
Eastern garter snake	<i>Thamnophis sirtalis</i>	none	X	X	X	X	C / N,F	38 40
eastern milk snake	<i>Lampropettis d. triangulum</i>	none	X	X	X	X	C / N,F	38 39
Fowler's toad	<i>Bufo woodhousei fowleri</i>	none	X	X	X	X	C / F	33 37

KEY:

Frequency:

A- abundant
C- common
R- rare
N- not expected

Activity:

N- nesting
H- hunting
R- resting
F- foraging

PROJECTION OF WILDLIFE ECOLOGICAL RESPONSE (POWER)

NELSON, POPE & VOORHIS, LLC, MICROCOMPUTER MODEL

SPECIES ADAPTABILITY

This portion of the appendix has been included to present the results of a computer program to identify "Species Adaptability." This list is another component of the program developed for use by Nelson, Pope & Voorhis, LLC to determine potential impacts of the proposed project on wildlife. In this application the "Adaptability" of the observed and expected species are shown. The "adaptability" as indicated in the table, refers to whether an individual species may potentially benefit from (+) a habitat change from natural to urban/suburban setting; or, be impacted (-), or remain constant (=), as a result of this change. These values are not intended to represent the dynamics of actual species on the subject site under post-development conditions. The column entitled "Comments" provides relevant information which was obtained from the literature, as regards special habits of the particular species, such as adaptability, nesting, food, etc. This column is particularly important in assessing the potential impacts to the species as a result of the proposed project. The preceding text considers the site specific aspects of the proposed development in regard to individual species. This appendix is included to provide the reader with the benefit of what the literature which was consulted in connection with the Habitat Suitability Model suggests, in terms of generalized species dynamics resulting from land use. References are the same as those cited in the previous Species List portion of the appendix.

Pine Barrens Species - Adaptability and Comments

Common Name	Scientific Name	Adapt.	Comments	References
Birds				
common bobwhite	<i>Colinus virginianus</i>	-	somewhat tolerant of humans during spring and summer months	4 8
gray catbird	<i>Dumetella carolinensis</i>	=	abundant around parks, urban and suburban areas	4 9
black capped chickadee	<i>Parus atricapillus</i>	=	abundant around parks, urban and suburban areas	4 11
American crow	<i>Corvus brachyrhynchos</i>	=	extremely adaptable; omnivorous	4 11
house finch	<i>Carpodacus mexicanus</i>	+	nests almost entirely on buildings; considered a pest species	4 20
purple finch	<i>Carpodacus purpureus</i>	-	inhabits parks, suburban areas, and coniferous forests	4 20
common flicker	<i>Colaptes auratus</i>	=	abundant around parks, suburban and urban areas	4 14
common grackle	<i>Quiscalus quiscula</i>	=	adapts well to urban and suburban habitats	4 6
ruffed grouse	<i>Bonasa umbellus</i>	-	prefers dense cover, thick woods; avoids humans	4 8
broad-winged hawk	<i>Buteo platypterus</i>	-	avoids humans; nests only in dense forests; prefers to be near water	4 16
Cooper's hawk	<i>Accipiter cooperii</i>	-	needs extensive woodland	4 17
red-tailed hawk	<i>Buteo jamaicensis</i>	-	needs 100 foot radius undisturbed area for nest	4 16
sharp-shinned hawk	<i>Accipiter striatus</i>	-	avoids humans; nests in heavily forested areas	4 16
blue jay	<i>Cyanocitta cristata</i>	=	extremely adaptable to human activity and other stresses	4 10
Northern (dark-eyed) junco	<i>Junco hyemalis</i>	-	prefers forested area with elevation >300 meters; no LI atlas record	4 21
Eastern kingbird	<i>Tyrannus tyrannus</i>	=	very adaptable to human activities; prefers open areas	4 15
golden-crowned kinglet	<i>Regulus satrapa</i>	-	prefers spruce vegetation; no atlas sightings on Long Island	4 7
ruby-crowned kinglet	<i>Regulus calendula</i>	-	occurs as non-breeding species; present during migration	4 7
white-breasted nuthatch	<i>Sitta carolinensis</i>	=	abundant in parks, urban and suburban areas	4 9
ovenbird	<i>Seiurus aurocapillus</i>	-	prefers open forest floor and woodlot greater than 35 acres	4 19
great-horned owl	<i>Bubo virginianus</i>	-	nocturnal; rare in wooded areas of less than 20 acres	4 17
long-eared owl	<i>Asio otus</i>	-	nocturnal; prefers dense forested areas near water	4 17
yellow-bellied sapsucker	<i>Sphyrapicus varius</i>	=	nests in tree cavity; found in parks, yards and gardens	14
pine siskin	<i>Carduelis pinus</i>	=	one atlas confirmed breeding record on Long Island	4 20
chipping sparrow	<i>Spizella passerina</i>	+	abundant around man made structures	4 21
fox sparrow	<i>Passerella iliaca</i>	-	boreal species, winters here in edge, thickets, brushy areas	20 21
song sparrow	<i>Melospiza melodia</i>	=	common to most habitats except deep forest, open field and marsh	4 22
European starling	<i>Sturnus vulgaris</i>	+	extremely adaptable to human activity; considered a pest	4 23
chimney swift	<i>Chaetura pelagica</i>	+	nests in chimneys, with few exceptions	4 42
scarlet tanager	<i>Piranga olivacea</i>	-	rare in wooded area of less than 50 acres; affected by fragmentation	4
brown thrasher	<i>Toxostoma rufum</i>	=	common in parks, suburban areas, wooded edges, dry open areas	4 9
hermit thrush	<i>Catharus guttatus</i>	=	not common on Long Island; when present, prefers pine barrens	4 7
wood thrush	<i>Hylocichla mustelina</i>	=	prefers vacant wood (trees >40 feet); may adapt of wooded suburban	4 7

Common Name	Scientific Name	Adapt.	Comments	References
pine warbler	<i>Dendroica pinus</i>	-	prefers pine forest; may appear in overgrown field	4 19
prairie warbler	<i>Dendroica discolor</i>	-	prefers scrub fields and open pine barrens habitat	4 19
yellow-rumped warbler	<i>Dendroica coronata</i>	-	prefers mixed and conifer forest; may be in yards	4 8
cedar waxwing	<i>Bombycilla cedrorum</i>	+	prefers open woodlands, orchards and residential areas	4 23 32
whip-poor-will	<i>Caprimulgus vociferous</i>	-	nocturnal; prefers open woods with adjacent fields	4 12
Eastern wood-peewee	<i>Contopus virens</i>	=	prefers suburban areas, parks and villages with shade trees	4 15
red-bellied woodpecker	<i>Melanerpes carolinus</i>	=	prefers forest openings; mostly found on Long Island north shore	4 14
house wren	<i>Troglodytes aedon</i>	=	found in suburban areas and gardens; nests in crevices of buildings	4 9
Mammals				
big-brown bat	<i>Eptesicus fuscus</i>	+	roosts in structures; found throughout LI; hunts over water	1 29
hoary bat	<i>Lasiurus borealis</i>	=	roosts in trees, sometimes found in parks	45
Keen's bat	<i>Myotis keenii</i>	+	roosts in buildings, crevices and bark; more common on eastern LI	1 29
little-brown bat	<i>Myotis lucifugus</i>	+	roosts in buildings and man made structures; hunts over water	1 29
red bat	<i>Lasiurus borealis</i>	-	feeds in marsh area; nests within 1000 yards of marsh in trees	1 29
Eastern pipistrelle	<i>Pipistrellus subflavus</i>	=	found near water in open woods, also found in buildings	1 29
silver-haired bat	<i>Lasionycteris noctivagans</i>	-	prefers wooded areas near water, primarily during summer months	1 29
Eastern chipmunk	<i>Tamias striatus</i>	=	prefers open woods, thickets, and rocky areas	1 29
Eastern cottontail	<i>Sylvilagus floridanus</i>	=	will adapt to suburban areas, if there is sufficient cover	1 29
white-tailed deer	<i>Odocoileus virginianus</i>	-	requires range of one-half square mile	1 25 29
red fox	<i>Vulpes vulpes</i>	-	builds den in wooded areas with loose-sandy soil and good drainage	1 29
Eastern mole	<i>Scalopus aquaticus</i>	=	tunnels underground	1 29
house mouse	<i>Mus musculus</i>	+	lives in association with man, not expected away from buildings	1 29
meadow-jumping mouse	<i>Zapus hudsonicus</i>	=	found around water in pine barrens; prefers open areas with grasses	1 29
white-footed mouse	<i>Peromyscus leucopus</i>	=	common to most all habitats; does not adapt well to human activity	1 29
Virginia opossum	<i>Didelphis virginiana</i>	=	common in suburban areas, woods, marsh and coastal areas	1 29
raccoon	<i>Procyon lotor</i>	+	nocturnal; very adaptive; found in urban and forest areas	1 29
masked shrew	<i>Sorex cinereus</i>	=	tunnels underground; common in wood and wet habitats	1 29
short-tailed shrew	<i>Blarina breuicauda</i>	=	tunnels underground; abundant in a variety of habitats	1 29
striped skunk	<i>Mephitis mephitis</i>	=	prefers mixed wood & brush within 2 miles of water; not expected on LI	1 29
Eastern gray squirrel	<i>Sciurus carolinensis</i>	=	found in parks, urban and suburban areas; very adaptable	1 29
southern-flying squirrel	<i>Glaucimys volans</i>	-	common in deep mixed, deciduous and coniferous woods	1 29
pine vole	<i>Microtus pinetorum</i>	=	tunnels underground; prefers sandy soil in woods and field; can swim	1 29
long-tailed weasel	<i>Mustela frenata</i>	-	prefers dense wood, but may appear in all land habitats near water	1 29
woodchuck	<i>Marmota monax</i>	-	appears primarily in scrub woods and brushy areas; not common on LI	1 29
Herptiles				
common gray treefrog	<i>Hyla versicolor</i>	-	prefer mossy trees near ponds	33 37
red-backed salamander	<i>Plethodon cinereus cinereus</i>	-	terrestrial, prevalent in moist situations	34 36

Common Name	Scientific Name	Adapt.	Comments	References
Eastern tiger salamander	Ambystoma tigrinum tigrinum	-	needs fishless pond or vernal pond with 500' vacant radius to breed	36 38
marbled salamander	Ambystoma opacum	=	moist to sandy areas; lays eggs in fall in low spots wet by rain	34 36 38
Eastern garter snake	Thamnophis sirtalis	=	occupies a variety of habitats	38 40
eastern hognose snake	Heterodon platyrhinos	=	sandy soil and sunny roadside; feeds on herptiles and insects	38
eastern milk snake	Lampropettis d. triangulum	=	occupies a variety of habitats	38 39
Eastern spadefoot toad	Scaphiopus holbrookii	-	nocturnal; burrows in sandy soil; eats insects, worms; gardens, etc.	33
Fowler's toad	Bufo woodhousei fowleri	-	found in suburban areas, gardens; breeds in shallow permanent ponds	33 37
Eastern box turtle	Terrepene carolina	-	terrestrial based species	41

Successional Woodland Species - Adaptability and Comments

Common Name	Scientific Name	Adapt.	Comments	References
Birds				
American crow	Corvus brachyrhynchos	=	extremely adaptable; omnivorous	4 11
American kestrel	Falco sparverius	-	adaptable; prefers open areas and parks; will nest near humans	4 17
American redstart	Setophaga ruticilla	-	urbanization and agriculture have negative effects	4 19
American robin	Turdus migratorius	=	very adaptable; abundant in parks; nests in man-made structures	4 7
American woodcock	Philhela minor	-	prefers moist woodland and thicket near open fields	4 30
barn swallow	Hirundo rustica	+	nests almost entirely on buildings	4 15
black capped chickadee	Parus atricapillus	=	abundant around parks, urban and suburban areas	4 11
black-and-white warbler	Mniotilta varia	-	builds nests under shrubs and/or trees	4 18
blue jay	Cyanocitta cristata	=	extremely adaptable to human activity and other stresses	4 10
blue-winged warbler	Vermivora pinus	-	primarily abandoned and overgrown field, and thickets	4 14
brown creeper	Certhia familiaris	-	prefers predominantly deciduous wooded areas	4 9
brown thrasher	Toxostoma rufum	=	common in parks,suburban areas, wooded edges, dry open areas	4 9
brown-headed cowbird	Molothrus ater	=	lays eggs in other bird's nests; some stay during winter	4 6
Carolina wren	Thryothorus ludovicianus	=	associated with woodland thickets and brushy areas, often near water	4 9
cedar waxwing	Bombycilla cedrorum	+	prefers open woodlands, orchards and residential areas	4 23 32
chestnut-sided warbler	Dendroica pensylvanica	-	prefers first growth woods, with some open brush area	4 19
common flicker	Colaptes auratus	=	abundant around parks, suburban and urban areas	4 14
common grackle	Quiscalus quiscula	=	adapts well to urban and suburban habitats	4 6
Cooper's hawk	Accipiter cooperii	-	needs extensive woodland	4 17
downy woodpecker	Picoides pubescens	=	found in parks and suburban areas	4 14
Eastern kingbird	Tyrannus tyrannus	=	very adaptable to human activities; prefers open areas	4 15
Eastern wood-peewee	Contopus virens	=	prefers suburban areas, parks and villages with shade trees	4 15
European starling	Sturnus vulgaris	+	extremely adaptable to human activity; considered a pest	4 23
fox sparrow	Passerella iliaca	-	boreal species, winters here in edge, thickets, brushy areas	20 21
golden-crowned kinglet	Regulus satrapa	-	prefers spruce vegetation; no atlas sightings on Long Island	4 7
gray catbird	Dumetella carolinensis	=	abundant around parks, urban and suburban areas	4 9
great-horned owl	Bubo virginianus	-	nocturnal; rare in wooded areas of less than 20 acres	4 17
hairy woodpecker	Picoides villosus	=	found mainly in deciduous forests	4 14
hermit thrush	Catharus guttatus	=	not common on Long Island; when present, prefers pine barrens	4 7
house finch	Carpodacus mexicanus	+	nests almost entirely on buildings; considered a pest species	4 20
house sparrow	Passer domesticus	+	prefers buildings, urban, suburban, gardens; considered a pest	4 20
house wren	Troglodytes aedon	=	found in suburban areas and gardens; nests in crevices of buildings	4 9
long-eared owl	Asio otus	-	nocturnal; prefers dense forested areas near water	4 17

Common Name	Scientific Name	Adapt.	Comments	References
mourning dove	Zenaida macroura	=	abundant around parks, urban and suburban areas	4 8
Northern (dark-eyed) junco	Junco hyemalis	-	prefers forested area with elevation >300 meters; no LI atlas record	4 21
Northern mockingbird	Mimus polyglottos	+	prefers to nest near humans	4 9
red-bellied woodpecker	Melanerpes carolinus	=	prefers forest openings; mostly found on Long Island north shore	4 14
red-eyed vireo	Vireo olivaceus	=	found in parks and suburban areas with shade trees and undergrowth	4 23
red-tailed hawk	Buteo jamaicensis	-	needs 100 foot radius undisturbed area for nest	4 16
rock dove	Columba livia	+	nests almost entirely on buildings; considered a pest species	4 8
rose-breasted grosbeak	Pheucticus ludovicianus	=	mainly found on north shore	4 20
ruby-crowned kinglet	Regulus calendula	-	occurs as non-breeding species; present during migration	4 7
ruffed grouse	Bonasa umbellus	-	prefers dense cover, thick woods; avoids humans	4 8
rufous-sided towhee	Pipilo erythrophthalmus	-	may be present year round on Long Island	4 20
sharp-shinned hawk	Accipiter striatus	-	avoids humans; nests in heavily forested areas	4 16
song sparrow	Melospiza melodia	=	common to most habitats except deep forest, open field and marsh	4 22
whip-poor-will	Caprimulgus vociferous	-	nocturnal; prefers open woods with adjacent fields	4 12
white-throated sparrow	Zonotrichia albicollis	-	prefers brushy areas and thick undergrowth	4 22
wood thrush	Hylocichla mustelina	=	prefers vacant wood (trees >40 feet); may adapt of wooded suburban	4 7
yellow-bellied sapsucker	Sphyrapicus varius	=	nests in tree cavity; found in parks, yards and gardens	14
yellow-billed cuckoo	Coccyzus americanus	-	avoids heavy urban areas; prefers wooded open or edges for nests	4 12
Mammals				
big-brown bat	Eptesicus fuscus	+	roosts in structures; found throughout LI; hunts over water	1 29
hoary bat	Lasiurus borealis	=	roosts in trees, sometimes found in parks	45
Keen's bat	Myotis keenii	+	roosts in buildings, crevices and bark; more common on eastern LI	1 29
little-brown bat	Myotis lucifugus	+	roosts in buildings and man made structures; hunts over water	1 29
red bat	Lasiurus borealis	-	feeds in marsh area; nests within 1000 yards of marsh in trees	1 29
silver-haired bat	Lasionycteris noctivagans	-	prefers wooded areas near water, primarily during summer months	1 29
Eastern chipmunk	Tamias striatus	=	prefers open woods, thickets, and rocky areas	1 29
Eastern cottontail	Sylvilagus floridanus	=	will adapt to suburban areas, if there is sufficient cover	1 29
white-tailed deer	Odocoileus virginianus	-	requires range of one-half square mile	1 25 29
red fox	Vulpes vulpes	-	builds den in wooded areas with loose-sandy soil and good drainage	1 29
Eastern mole	Scalopus aquaticus	=	tunnels underground	1 29
meadow-jumping mouse	Zapus hudsonicus	=	found around water in pine barrens; prefers open areas with grasses	1 29
white-footed mouse	Peromyscus leucopus	=	common to most all habitats; does not adapt well to human activity	1 29
Virginia opossum	Didelphis virginiana	=	common in suburban areas, woods, marsh and coastal areas	1 29
raccoon	Procyon lotor	+	nocturnal; very adaptive; found in urban and forest areas	1 29
masked shrew	Sorex cinereus	=	tunnels underground; common in wood and wet habitats	1 29
short-tailed shrew	Blarina breuicauda	=	tunnels underground; abundant in a variety of habitats	1 29

Common Name	Scientific Name	Adapt.	Comments	References
striped skunk	Mephitis mephitis	=	prefers mixed wood & brush within 2 miles of water; not expected on LI	1 29
Eastern gray squirrel	Sciurus carolinensis	=	found in parks, urban and suburban areas; very adaptable	1 29
meadow vole	Microtus pennsylvanicus	=	tunnels underground; prefers open woodland	29 45
pine vole	Microtus pinetorum	=	tunnels underground; prefers sandy soil in woods and field; can swim	1 29
long-tailed weasel	Mustela frenata	-	prefers dense wood, but may appear in all land habitats near water	1 29
woodchuck	Marmota monax	-	appears primarily in scrub woods and brushy areas; not common on LI	1 29
Herptiles				
Eastern garter snake	Thamnophis sirtalis	=	occupies a variety of habitats	38 40
eastern hognose snake	Heterodon platyrhinos	=	sandy soil and sunny roadside; feeds on herptiles and insects	38
eastern milk snake	Lampropeltis d. triangulum	=	occupies a variety of habitats	38 39

Successional Shrubland Species - Adaptability and Comments

Common Name	Scientific Name	Adapt.	Comments	References
Birds				
red-winged blackbird	Agelaius phoeniceus	=	needs water	4 6
common bobwhite	Colinus virginianus	-	somewhat tolerant of humans during spring and summer months	4 8
indigo bunting	Passerina cyanea	-	inhabits open woodlands with dense thickets for cover	4 20
Northern cardinal	Cardinalis cardinalis	=	found around gardens, yards, parks	4 20
gray catbird	Dumetella carolinensis	=	abundant around parks, urban and suburban areas	4 9
black capped chickadee	Parus atricapillus	=	abundant around parks, urban and suburban areas	4 11
brown-headed cowbird	Molothrus ater	=	lays eggs in other bird's nests; some stay during winter	4 6
American crow	Corvus brachyrhynchos	=	extremely adaptable; omnivorous	4 11
black-billed cuckoo	Coccyzus erythrophthalmus	-	avoids human activities	4 11
yellow-billed cuckoo	Coccyzus americanus	-	avoids heavy urban areas; prefers wooded open or edges for nests	4 12
mourning dove	Zenaida macroura	=	abundant around parks, urban and suburban areas	4 8
American goldfinch	Carduelis tristis	=	prefers diet of thistles and dandelions	4 20
common flicker	Colaptes auratus	=	abundant around parks, suburban and urban areas	4 14
least flycatcher	Empidonax minimus	=	prefers open areas and woodland borders; uncommon on Long Island	4 15
willow flycatcher	Empidonax traillii	-	found mostly on south shore and western north shore areas	4 15
common grackle	Quiscalus quiscula	=	adapts well to urban and suburban habitats	4 6
ruffed grouse	Bonasa umbellus	-	prefers dense cover, thick woods; avoids humans	4 8
rose-breasted grosbeak	Pheucticus ludovicianus	=	mainly found on north shore	4 20
Northern harrier	Circus cyaneus	-	avoids humans; extremely protective of nests	4 16
Cooper's hawk	Accipiter cooperii	-	needs extensive woodland	4 17
red-tailed hawk	Buteo jamaicensis	-	needs 100 foot radius undisturbed area for nest	4 16
sharp-shinned hawk	Accipiter striatus	-	avoids humans; nests in heavily forested areas	4 16
blue jay	Cyanocitta cristata	=	extremely adaptable to human activity and other stresses	4 10
Northern (dark-eyed) junco	Junco hyemalis	-	prefers forested area with elevation >300 meters; no LI atlas record	4 21
American kestrel	Falco sparverius	-	adaptable; prefers open areas and parks; will nest near humans	4 17
Eastern kingbird	Tyrannus tyrannus	=	very adaptable to human activities; prefers open areas	4 15
golden-crowned kinglet	Regulus satrapa	-	prefers spruce vegetation; no atlas sightings on Long Island	4 7
ruby-crowned kinglet	Regulus calendula	-	occurs as non-breeding species; present during migration	4 7
Eastern meadowlark	Sturnella magna	-	found in marshes during winter months	4 6
Northern mockingbird	Mimus polyglottos	+	prefers to nest near humans	4 9
barn owl	Tyto alba	=	hunts in open areas, nests in man made structures and hollow trees	4 17
great-horned owl	Bubo virginianus	-	nocturnal; rare in wooded areas of less than 20 acres	4 17
ring-necked pheasant	Phasianus colchicus	-	needs fields with cover along edge	4 8

Common Name	Scientific Name	Adapt.	Comments	References
American redstart	Setophaga ruticilla	-	urbanization and agriculture have negative effects	4 19
American robin	Turdus migratorius	=	very adaptable; abundant in parks; nests in man-made structures	4 7
pine siskin	Carduelis pinus	=	one atlas confirmed breeding record on Long Island	4 20
fox sparrow	Passerella iliaca	-	boreal species, winters here in edge, thickets, brushy areas	20 21
field sparrow	Spizella pusilla	-	associated with grasslands, fields and brushy wooded edges	4 21
grasshopper sparrow	Ammodramus savannarum	-	requires grasslands	4 20
song sparrow	Melospiza melodia	=	common to most habitats except deep forest, open field and marsh	4 22
swamp sparrow	Melospiza georgiana	-	prefers fresh water marshes; may be found in weedy fields, parks	4 22
white-crowned sparrow	Zonotrichia leucophrys	=	often found in suburban areas and city parks	22 32
white-throated sparrow	Zonotrichia albicollis	-	prefers brushy areas and thick undergrowth	4 22
European starling	Sturnus vulgaris	+	extremely adaptable to human activity; considered a pest	4 23
barn swallow	Hirundo rustica	+	nests almost entirely on buildings	4 15
brown thrasher	Toxostoma rufum	=	common in parks and suburban areas, wooded edges and dry open areas	4 9
hermit thrush	Catharus guttatus	=	not common on Long Island; when present, prefers pine barrens	4 7
wood thrush	Hylocichla mustelina	=	prefers vacant wood (trees >40 feet); may adapt of wooded suburban	4 7
rufous-sided towhee	Pipilo erythrophthalmus	-	may be present year round on Long Island	4 20
white eyed vireo	Vireo griseus	-	avoids human activity; prefers dense swampy thickets	4 23
black-and-white warbler	Mniotilta varia	-	builds nests under shrubs and/or trees	4 18
blue-winged warbler	Vermivora pinus	-	primarily abandoned and overgrown field, and thickets	4 14
chestnut-sided warbler	Dendroica pensylvanica	-	prefers first growth woods, with some open brush area	4 19
prairie warbler	Dendroica discolor	-	prefers scrub fields and open pine barrens habitat	4 19
cedar waxwing	Bombycilla cedrorum	+	prefers open woodlands, orchards and residential areas	4 23 32
whip-poor-will	Caprimulgus vociferous	-	nocturnal; prefers open woods with adjacent fields	4 12
American woodcock	Philhela minor	-	prefers moist woodland and thicket near open fields	4 30
Carolina wren	Thryothorus ludovicianus	=	associated with woodland thickets and brushy areas, often near water	4 9
house wren	Troglodytes aedon	=	found in suburban areas and gardens; nests in crevices of buildings	4 9
common yellowthroat	Geothlypis trichas	=	found in all open brushy wet areas	4 19
Mammals				
Eastern chipmunk	Tamias striatus	=	prefers open woods, thickets, and rocky areas	1 29
Eastern cottontail	Sylvilagus floridanus	=	will adapt to suburban areas, if there is sufficient cover	1 29
white-tailed deer	Odocoileus virginianus	-	requires range of one-half square mile	1 25 29
red fox	Vulpes vulpes	-	builds den in wooded areas with loose-sandy soil and good drainage	1 29
Eastern mole	Scalopus aquaticus	=	tunnels underground	1 29
house mouse	Mus musculus	+	lives in association with man, not expected away from buildings	1 29
meadow-jumping mouse	Zapus hudsonicus	=	found around water in pine barrens; prefers open areas with grasses	1 29
white-footed mouse	Peromyscus leucopus	=	common to most all habitats; does not adapt well to human activity	1 29
Virginia opossum	Didelphis virginiana	=	common in suburban areas, as well as woods, marsh and coastal areas	1 29

Common Name	Scientific Name	Adapt.	Comments	References
racoon	<i>Procyon lotor</i>	+	nocturnal; very adaptive; found in urban and forest areas	1 29
black rat	<i>Rattus rattus</i>	=	lives in association with man, mainly city water front buildings	1 29
short-tailed shrew	<i>Blarina breuicauda</i>	=	tunnels underground; abundant in a variety of habitats	1 29
striped skunk	<i>Mephitis mephitis</i>	=	prefers mixed wood & brush within 2 miles of water; not expected on LI	1 29
meadow vole	<i>Microtus pennsylvanicus</i>	=	tunnels underground; prefers open woodland	29 45
pine vole	<i>Microtus pinetorum</i>	=	tunnels underground; prefers sandy soil in woods and field; can swim	1 29
long-tailed weasel	<i>Mustela frenata</i>	-	prefers dense wood, but may appear in all land habitats near water	1 29
woodchuck	<i>Marmota monax</i>	-	appears primarily in scrub woods and brushy areas; not common on LI	1 29
Herptiles				
Eastern garter snake	<i>Thamnophis sirtalis</i>	=	occupies a variety of habitats	38 40
eastern hognose snake	<i>Heterodon platyrhinos</i>	=	sandy soil and sunny roadside; feeds on herptiles and insects	38
eastern milk snake	<i>Lampropeltis d. triangulum</i>	=	occupies a variety of habitats	38 39
Fowler's toad	<i>Bufo woodhousei fowleri</i>	-	found in suburban areas, gardens; breeds in shallow permanent ponds	33 37

Successional Old Field Species - Adaptability and Comments

Common Name	Scientific Name	Adapt.	Comments	References
Birds				
American crow	<i>Corvus brachyrhynchos</i>	=	extremely adaptable; omnivorous	4 11
American goldfinch	<i>Carduelis tristis</i>	=	prefers diet of thistles and dandelions	4 20
American kestrel	<i>Falco sparverius</i>	-	adaptable; prefers open areas and parks; will nest near humans	4 17
American robin	<i>Turdus migratorius</i>	=	very adaptable; abundant in parks; nests in man-made structures	4 7
American woodcock	<i>Philhela minor</i>	-	prefers moist woodland and thicket near open fields	4 30
barn owl	<i>Tyto alba</i>	=	hunts in open areas, nests in man made structures and hollow trees	4 17
barn swallow	<i>Hirundo rustica</i>	+	nests almost entirely on buildings	4 15
black capped chickadee	<i>Parus atricapillus</i>	=	abundant around parks, urban and suburban areas	4 11
black-and-white warbler	<i>Mniotilta varia</i>	-	builds nests under shrubs and/or trees	4 18
black-billed cuckoo	<i>Coccyzus erythrophthalmus</i>	-	avoids human activities	4 11
blue jay	<i>Cyanocitta cristata</i>	=	extremely adaptable to human activity and other stresses	4 10
blue-winged warbler	<i>Vermivora pinus</i>	-	primarily abandoned and overgrown field, and thickets	4 14
brown thrasher	<i>Toxostoma rufum</i>	=	common in parks and suburban areas, wooded edges and dry open areas	4 9
brown-headed cowbird	<i>Molothrus ater</i>	=	lays eggs in other bird's nests; some stay during winter	4 6
cedar waxwing	<i>Bombycilla cedrorum</i>	+	prefers open woodlands, orchards and residential areas	4 23 32
chestnut-sided warbler	<i>Dendroica pensylvanica</i>	-	prefers first growth woods, with some open brush area	4 19
chipping sparrow	<i>Spizella passerina</i>	+	abundant around man made structures	4 21
common bobwhite	<i>Colinus virginianus</i>	-	somewhat tolerant of humans during spring and summer months	4 8
common flicker	<i>Colaptes auratus</i>	=	abundant around parks, suburban and urban areas	4 14
common grackle	<i>Quiscalus quiscula</i>	=	adapts well to urban and suburban habitats	4 6
common nighthawk	<i>Chordeiles minor</i>	=	primarily a grassland specie; will nest in burnt areas and roofs	4 12
common yellowthroat	<i>Geothlypis trichas</i>	=	found in all open brushy wet areas	4 19
Cooper's hawk	<i>Accipiter cooperii</i>	-	needs extensive woodland	4 17
Eastern bluebird	<i>Sialia sialis</i>	-	found almost entirely in nesting boxes, extremely rare in wild	4 7
Eastern kingbird	<i>Tyrannus tyrannus</i>	=	very adaptable to human activities; prefers open areas	4 15
Eastern meadowlark	<i>Sturnella magna</i>	-	found in marshes during winter months	4 6
European starling	<i>Sturnus vulgaris</i>	+	extremely adaptable to human activity; considered a pest	4 23
field sparrow	<i>Spizella pusilla</i>	-	associated with grasslands, fields and brushy wooded edges	4 21
fox sparrow	<i>Passerella iliaca</i>	-	boreal species, winters here in edge, thickets, brushy areas	20 21
grasshopper sparrow	<i>Ammodramus savannarum</i>	-	requires grasslands	4 20
gray catbird	<i>Dumetella carolinensis</i>	=	abundant around parks, urban and suburban areas	4 9
great-horned owl	<i>Bubo virginianus</i>	-	nocturnal; rare in wooded areas of less than 20 acres	4 17
house finch	<i>Carpodacus mexicanus</i>	+	nests almost entirely on buildings; considered a pest species	4 20

Common Name	Scientific Name	Adapt.	Comments	References
house sparrow	<i>Passer domesticus</i>	+	prefers buildings, urban, suburban, gardens; considered a pest	4 20
house wren	<i>Troglodytes aedon</i>	=	found in suburban areas and gardens; nests in crevices of buildings	4 9
indigo bunting	<i>Passerina cyanea</i>	-	inhabits open woodlands with dense thickets for cover	4 20
least flycatcher	<i>Empidonax minimus</i>	=	prefers open areas and woodland borders; uncommon on Long Island	4 15
mourning dove	<i>Zenaida macroura</i>	=	abundant around parks, urban and suburban areas	4 8
Northern (dark-eyed) junco	<i>Junco hyemalis</i>	-	prefers forested area with elevation >300 meters; no LI atlas record	4 21
Northern cardinal	<i>Cardinalis cardinalis</i>	=	found around gardens, yards, parks	4 20
Northern harrier	<i>Circus cyaneus</i>	-	avoids humans; extremely protective of nests	4 16
Northern mockingbird	<i>Mimus polyglottos</i>	+	prefers to nest near humans	4 9
prairie warbler	<i>Dendroica discolor</i>	-	prefers scrub fields and open pine barrens habitat	4 19
red-headed woodpecker	<i>Melanerpes erythrocephalus</i>	=	prefers open woodlands, parks and suburban areas	4 14
red-tailed hawk	<i>Buteo jamaicensis</i>	-	needs 100 foot radius undisturbed area for nest	4 16
red-winged blackbird	<i>Agelaius phoeniceus</i>	=	needs water	4 6
ring-necked pheasant	<i>Phasianus colchicus</i>	-	needs fields with cover along edge	4 8
rock dove	<i>Columba livia</i>	+	nests almost entirely on buildings; considered a pest species	4 8
rose-breasted grosbeak	<i>Pheucticus ludovicianus</i>	=	mainly found on north shore	4 20
ruffed grouse	<i>Bonasa umbellus</i>	-	prefers dense cover, thick woods; avoids humans	4 8
rufous-sided towhee	<i>Pipilo erythrophthalmus</i>	-	may be present year round on Long Island	4 20
Savannah sparrow	<i>Passerculus sandwichensis</i>	-	found in shore areas; not expected inland	4 21
sharp-shinned hawk	<i>Accipiter striatus</i>	-	avoids humans; nests in heavily forested areas	4 16
song sparrow	<i>Melospiza melodia</i>	=	common to most habitats except deep forest, open field and marsh	4 22
swamp sparrow	<i>Melospiza georgiana</i>	-	prefers fresh water marshes; may be found in weedy fields, parks	4 22
whip-poor-will	<i>Caprimulgus vociferous</i>	-	nocturnal; prefers open woods with adjacent fields	4 12
white-crowned sparrow	<i>Zonotrichia leucophrys</i>	=	often found in suburban areas and city parks	22 32
willow flycatcher	<i>Empidonax traillii</i>	-	found mostly on south shore and western north shore areas	4 15
yellow warbler	<i>Dendrocica petchia</i>	=	rare breeder on LI, winter sps, abundant in parks & yards	4 18
yellow-billed cuckoo	<i>Coccyzus americanus</i>	-	avoids heavy urban areas; prefers wooded open or edges for nests	4 12
Mammals				
Eastern chipmunk	<i>Tamias striatus</i>	=	prefers open woods, thickets, and rocky areas	1 29
Eastern cottontail	<i>Sylvilagus floridanus</i>	=	will adapt to suburban areas, if there is sufficient cover	1 29
white-tailed deer	<i>Odocoileus virginianus</i>	-	requires range of one-half square mile	1 25 29
red fox	<i>Vulpes vulpes</i>	-	builds den in wooded areas with loose-sandy soil and good drainage	1 29
Eastern mole	<i>Scalopus aquaticus</i>	=	tunnels underground	1 29
house mouse	<i>Mus musculus</i>	+	lives in association with man, not expected away from buildings	1 29
meadow-jumping mouse	<i>Zapus hudsonicus</i>	=	found around water in pine barrens; prefers open areas with grasses	1 29
white-footed mouse	<i>Peromyscus leucopus</i>	=	common to most all habitats; does not adapt well to human activity	1 29
Virginia opossum	<i>Didelphis virginiana</i>	=	common in suburban areas, as well as woods, marsh and coastal areas	1 29

Common Name	Scientific Name	Adapt.	Comments	References
raccoon	<i>Procyon lotor</i>	+	nocturnal; very adaptive; found in urban and forest areas	1 29
black rat	<i>Rattus rattus</i>	=	lives in association with man, mainly city water front buildings	1 29
Norway rat	<i>Rattus norvegicus</i>	+	nocturnal; usually associated with human activity	1 29
least shrew	<i>Cryptotis parva</i>	-	not commonly documented on Long Island	1 29
short-tailed shrew	<i>Blarina breuicauda</i>	=	tunnels underground; abundant in a variety of habitats	1 29
striped skunk	<i>Mephitis mephitis</i>	=	prefers mixed wood & brush within 2 miles of water; not expected on LI	1 29
meadow vole	<i>Microtus pennsylvanicus</i>	=	tunnels underground; prefers open woodland	29 45
pine vole	<i>Microtus pinetorum</i>	=	tunnels underground; prefers sandy soil in woods and field; can swim	1 29
long-tailed weasel	<i>Mustela frenata</i>	-	prefers dense wood, but may appear in all land habitats near water	1 29
woodchuck	<i>Marmota monax</i>	-	appears primarily in scrub woods and brushy areas; not common on LI	1 29
Herptiles				
Eastern garter snake	<i>Thamnophis sirtalis</i>	=	occupies a variety of habitats	38 40
eastern hognose snake	<i>Heterodon platyrhinos</i>	=	sandy soil and sunny roadside; feeds on herptiles and insects	38
eastern milk snake	<i>Lampropeltis d. triangulum</i>	=	occupies a variety of habitats	38 39
Fowler's toad	<i>Bufo woodhousei fowleri</i>	-	found in suburban areas, gardens; breeds in shallow permanent ponds	33 37

Wooded Swamp Species : Adaptability and Comments

Common Name	Scientific Name	Adapt.	Comments	References
Birds				
red-winged blackbird	Agelaius phoeniceus	=	needs water	4 6
Northern cardinal	Cardinalis cardinalis	=	found around gardens, yards, parks	4 20
wood duck	Aix sponsa	-	prefers wooded rivers and ponds, and wooded swamps	4 27
great-crested flycatcher	Myiarchus crinitus	-	prefers deciduous forests and deciduous open woodland	4 15
blue-grey gnatcatcher	Poliophtila caerulea	=	prefers dense foliated trees along water ways	4 7
rose-breasted grosbeak	Pheucticus ludovicianus	=	mainly found on north shore	4 20
yellow-crowned night-heron	Nycticorax violaceus	-	nests in low coastal shrubbery; prefers islands	4 26
mallard	Anas platyrhynchos	-	adaptable to human activity	4 27
white-breasted nuthatch	Sitta carolinensis	=	abundant in parks, urban and suburban areas	4 9
osprey	Pandion haliaetus	-	associated with seacoast, sometimes lakes and rivers	4 16
great-horned owl	Bubo virginianus	-	nocturnal; rare in wooded areas of less than 20 acres	4 17
saw-whet owl	Aegolius acadicus	-	nocturnal; low moist coniferous; winter in parks, yards, thickets	4 17
Eastern phoebe	Sayornis phoebe	=	prefers open wood near stream; nests in log, building, bridge, cliff	4 15
spotted sandpiper	Actitis macularia	-	nests on ground in grassy areas	4 31 32
swamp sparrow	Melospiza georgiana	-	prefers fresh water marshes; may be found in weedy fields, parks	4 22
European starling	Sturnus vulgaris	+	extremely adaptable to human activity; considered a pest	4 23
tree swallow	Tachycineta bicolor	=	always nests near water	4 15
tufted titmouse	Parus bicolor	=	common in suburban areas	4 11
veery	Catharus fuscescens	-	prefers damp forest with undergrowth; affected by fragmentation	4 7
yellow warbler	Dendrocica petchia	=	rare breeder on LI, winter sps, abundant in parks & yards	4 18
American woodcock	Philhela minor	-	prefers moist woodland and thicket near open fields	4 30
downy woodpecker	Picoides pubescens	=	found in parks and suburban areas	4 14
hairy woodpecker	Picoides villosus	=	found mainly in deciduous forests	4 14
red-bellied woodpecker	Melanerpes carolinus	=	prefers forest openings; mostly found on Long Island north shore	4 14
Mammals				
big-brown bat	Eptesicus fuscus	+	roosts in structures; found throughout LI; hunts over water	1 29
little-brown bat	Myotis lucifugus	+	roosts in buildings and man made structures; hunts over water	1 29
Eastern cottontail	Sylvilagus floridanus	=	will adapt to suburban areas, if there is sufficient cover	1 29
white-tailed deer	Odocoileus virginianus	-	requires range of one-half square mile	1 25 29
red fox	Vulpes vulpes	-	builds den in wooded areas with loose-sandy soil and good drainage	1 29
mink	Mustela vison	-	prefers wetlands surrounded by forested areas	1
star-nosed mole	Condylura cristata	-	prefers wet, black, soil with earthworms	1 29
meadow-jumping mouse	Zapus hudsonicus	=	found around water in pine barrens; prefers open areas with grasses	1 29

Common Name	Scientific Name	Adapt.	Comments	References
white-footed mouse	<i>Peromyscus leucopus</i>	=	common to most all habitats; does not adapt well to human activity	1 29
muskrat	<i>Ondarta zibethicus</i>	-	prefers damp and marshy fresh and salt water habitats	1 29
Virginia opossum	<i>Didelphis virginiana</i>	=	common in suburban areas, woods, marsh and coastal areas	1 29
raccoon	<i>Procyon lotor</i>	+	nocturnal; very adaptive; found in urban and forest areas	1 29
Norway rat	<i>Rattus norvegicus</i>	+	nocturnal; usually associated with human activity	1 29
masked shrew	<i>Sorex cinereus</i>	=	tunnels underground; common in wood and wet habitats	1 29
short-tailed shrew	<i>Blarina breuicauda</i>	=	tunnels underground; abundant in a variety of habitats	1 29
Eastern gray squirrel	<i>Sciurus carolinensis</i>	=	found in parks, urban and suburban areas; very adaptable	1 29
southern-flying squirrel	<i>Glaucomys volans</i>	-	common in deep mixed, deciduous and coniferous woods	1 29
meadow vole	<i>Microtus pennsylvanicus</i>	=	tunnels underground; prefers open woodland	29 45
pine vole	<i>Microtus pinetorum</i>	=	tunnels underground; prefers sandy soil in woods and field; can swim	1 29
long-tailed weasel	<i>Mustela frenata</i>	-	prefers dense wood, but may appear in all land habitats near water	1 29
Herptiles				
bull frog	<i>Rana catesbeiana</i>	-	strictly aquatic, wooded lakes	33 34 35 37
common gray treefrog	<i>Hyla versicolor</i>	-	prefer mossy trees near ponds	33 37
green frog	<i>Rana clamitans</i>	-	mainly aquatic species	33 35 37
Southern leopard frog	<i>Rana pipiens sphenoccephala</i>	-	prefers shallow fresh to brackish pond; may be in meadow in summer	35 37 38
wood frog	<i>Rana sylvatica</i>	-	prefers leafy pools and transient pools in wooded areas	33 37
red-spotted newt	<i>Notophthalmus viridescens</i>	-	prefers shallow ponds in wooded areas; open moist woods	36 38
spring peeper	<i>Hyla crucifer</i>	=	prefers pools/marsh near woodland; found high in trees in summer	33 35 38
spotted salamander	<i>Ambystoma maculatum</i>	-	will breed in pond or vernal ponds in late March, early April	34 36 38
Eastern tiger salamander	<i>Ambystoma tigrinum tigrinum</i>	-	needs fishless pond or vernal pond with 500' vacant radius to breed	36 38
marbled salamander	<i>Ambystoma opacum</i>	=	moist to sandy areas; lays eggs in fall in low spots wet by rain	34 36 38
Eastern garter snake	<i>Thamnophis sirtalis</i>	=	occupies a variety of habitats	38 40
eastern milk snake	<i>Lampropettis d. triangulum</i>	=	occupies a variety of habitats	38 39
Eastern ribbon snake	<i>Thamnophis s. sauritus</i>	=	semi-aquatic specie seldom wanders far from wet areas	38 40
Northern brown snake	<i>Storeria dekayi</i>	=	prefers fresh marsh, moist woods, but, adapts to urban environment	38
Northern ringneck snake	<i>Diadophis punctatus</i>	=	prefers secluded moist areas under logs/stones; can adapt to suburb	38
Northern water snake	<i>Natrix sipedon sipedon</i>	-	common in swamp, bog, marsh, stream, pond and lake environments	38 39
painted turtle	<i>Chrysemys picta</i>	-	prefers small bodies of water	38

Terrestrial Cultural Species : Adaptability

Common Name	Scientific Name	Adapt.	Comments	References
Birds				
red-winged blackbird	Agelaius phoeniceus	=	needs water	4 6
Eastern bluebird	Sialia sialis	-	found almost entirely in nesting boxes, extremely rare in wild	4 7
common bobwhite	Colinus virginianus	-	somewhat tolerant of humans during spring and summer months	4 8
indigo bunting	Passerina cyanea	-	inhabits open woodlands with dense thickets for cover	4 20
Northern cardinal	Cardinalis cardinalis	=	found around gardens, yards, parks	4 20
gray catbird	Dumetella carolinensis	=	abundant around parks, urban and suburban areas	4 9
brown-headed cowbird	Molothrus ater	=	lays eggs in other bird's nests; some stay during winter	4 6
American crow	Corvus brachyrhynchos	=	extremely adaptable; omnivorous	4 11
mourning dove	Zenaida macroura	=	abundant around parks, urban and suburban areas	4 8
rock dove	Columba livia	+	nests almost entirely on buildings; considered a pest species	4 8
great-crested flycatcher	Myiarchus crinitus	-	prefers deciduous forests and deciduous open woodland	4 15
common grackle	Quiscalus quiscula	=	adapts well to urban and suburban habitats	4 6
Northern harrier	Circus cyaneus	-	avoids humans; extremely protective of nests	4 16
Cooper's hawk	Accipiter cooperii	-	needs extensive woodland	4 17
red-tailed hawk	Buteo jamaicensis	-	needs 100 foot radius undisturbed area for nest	4 16
sharp-shinned hawk	Accipiter striatus	-	avoids humans; nests in heavily forested areas	4 16
blue jay	Cyanocitta cristata	=	extremely adaptable to human activity and other stresses	4 10
American kestrel	Falco sparverius	-	adaptable; prefers open areas and parks; will nest near humans	4 17
killdeer	Charadrius vociferus	=	grassland species prefers coastal and wet areas with sparse growth	4 31 32
Eastern kingbird	Tyrannus tyrannus	=	very adaptable to human activities; prefers open areas	4 15
Eastern meadowlark	Sturnella magna	-	found in marshes during winter months	4 6
Northern mockingbird	Mimus polyglottos	+	prefers to nest near humans	4 9
common nighthawk	Chordeiles minor	=	primarily a grassland specie; will nest in burnt areas and roofs	4 12
barn owl	Tyto alba	=	hunts in open areas, nests in man made structures and hollow trees	4 17
great-horned owl	Bubo virginianus	-	nocturnal; rare in wooded areas of less than 20 acres	4 17
ring-necked pheasant	Phasianus colchicus	-	needs fields with cover along edge	4 8
black-bellied plover	Pluvialis squatarola	-	non-breeder on LI; may forage during summer, occasionally winter	31 32
American robin	Turdus migratorius	=	very adaptable; abundant in parks; nests in man-made structures	4 7
chipping sparrow	Spizella passerina	+	abundant around man made structures	4 21
field sparrow	Spizella pusilla	-	associated with grasslands, fields and brushy wooded edges	4 21
grasshopper sparrow	Ammodramus savannarum	-	requires grasslands	4 20
house sparrow	Passer domesticus	+	prefers buildings, urban, suburban, gardens; considered a pest	4 20
Savannah sparrow	Passerculus sandwichensis	-	found in shore areas; not expected inland	4 21

Common Name	Scientific Name	Adapt.	Comments	References
song sparrow	Melospiza melodia	=	common to most habitats except deep forest, open field and marsh	4 22
white-crowned sparrow	Zonotrichia leucophrys	=	often found in suburban areas and city parks	22 32
European starling	Sturnus vulgaris	+	extremely adaptable to human activity; considered a pest	4 23
barn swallow	Hirundo rustica	+	nests almost entirely on buildings	4 15
chimney swift	Chaetura pelagica	+	nests in chimneys, with few exceptions	4 42
brown thrasher	Toxostoma rufum	=	common in parks, suburban areas, wooded edges, dry open areas	4 9
chestnut-sided warbler	Dendroica pensylvanica	-	prefers first growth woods, with some open brush area	4 19
cedar waxwing	Bombycilla cedrorum	+	prefers open woodlands, orchards and residential areas	4 23 32
whip-poor-will	Caprimulgus vociferous	-	nocturnal; prefers open woods with adjacent fields	4 12
American woodcock	Philhela minor	-	prefers moist woodland and thicket near open fields	4 30
red-headed woodpecker	Melanerpes erythrocephalus	=	prefers open woodlands, parks and suburban areas	4 14
house wren	Troglodytes aedon	=	found in suburban areas and gardens; nests in crevices of buildings	4 9
Mammals				
Eastern pipistrelle	Pipistrellus subflavus	=	found near water in open woods, also found in buildings	1 29
Eastern cottontail	Sylvilagus floridanus	=	will adapt to suburban areas, if there is sufficient cover	1 29
white-tailed deer	Odocoileus virginianus	-	requires range of one-half square mile	1 25 29
red fox	Vulpes vulpes	-	builds den in wooded areas with loose-sandy soil and good drainage	1 29
house mouse	Mus musculus	+	lives in association with man, not expected away from buildings	1 29
meadow-jumping mouse	Zapus hudsonicus	=	found around water in pine barrens; prefers open areas with grasses	1 29
white-footed mouse	Peromyscus leucopus	=	common to most all habitats; does not adapt well to human activity	1 29
Virginia opossum	Didelphis virginiana	=	common in suburban areas, woods, marsh and coastal areas	1 29
raccoon	Procyon lotor	+	nocturnal; very adaptive; found in urban and forest areas	1 29
black rat	Rattus rattus	=	lives in association with man, mainly city water front buildings	1 29
Norway rat	Rattus norvegicus	+	nocturnal; usually associated with human activity	1 29
least shrew	Cryptotis parva	-	not commonly documented on Long Island	1 29
striped skunk	Mephitis mephitis	=	prefers mixed wood & brush within 2 miles of water; not expected on LI	1 29
pine vole	Microtus pinetorum	=	tunnels underground; prefers sandy soil in woods and field; can swim	1 29
long-tailed weasel	Mustela frenata	-	prefers dense wood, but may appear in all land habitats near water	1 29
Herptiles				
Eastern garter snake	Thamnophis sirtalis	=	occupies a variety of habitats	38 40
eastern milk snake	Lampropettis d. triangulum	=	occupies a variety of habitats	38 39
Fowler's toad	Bufo woodhousei fowleri	-	found in suburban areas, gardens; breeds in shallow permanent ponds	33 37

REFERENCES FOR WILDLIFE MATRIX

Refer.	Publication	Refer.	Publication
1	Connor, P.F. 1971. The Mammals of Long Island. NYS Museum Science Service Bulletin 416 SUNY, Albany.	9	Bent, A.C. 1964. Life Histories of North American Nuthatches, Wrens Dover Pub., NY.
4	Andrle, R.E., and J.R. Carroll. 1988. The Atlas Of Breeding Birds in New York State. Cornell University Press, Ithaca.	10	Bent, A.C. 1964. Life Histories of North American Jays, Crows, and Titmice, pt. 1. Dover Pub., NY
5	Pontin, A.J. 1982. Competition an Advanced Publishing Program, Boston, Massachusetts.	11	Bent, A.C. 1964. Life Histories of North American Jays, Crows, and Titmice, pt. 2. Dover Pub., NY
6	Bent, A.C. 1965. Life Histories of North American Black birds, Orioles, Tangers, and their allies. Dover Pub., NY.	12	Bent, A.C. 1964. Life Histories of North American Cuckoos, Goatsuckers, Hummingbirds, and their allies, pt. 1. Dover Pub., NY
7	Bent, A.C. 1964. Life Histories of North American Thrushes, Kinglets, and their allies. Dover Pub., NY.	13	Bent, A.C. 1964. Life Histories of North American Cuckoos, Goatsuckers, Hummingbirds, and their allies, pt. 2. Dover Pub., NY.
8	Bent, A.C. 1963. Life Histories of North American Gallinaceous Birds. Dover Pub., NY.	14	Bent, A.C. 1964. Life Histories of North American Woodpeckers. Dover Pub., NY.

- 15** Bent, A.C. 1963. Life Histories of North American Flycatchers, Larks, Swallows, and their allies. Dover Pub., NY.
- 16** Bent, A.C. 1961. Life Histories of North American Birds of Prey, pt 1. Dover Pub., NY.
- 17** Bent, A.C. 1961. Life Histories of North American Birds of Prey, pt 1. Dover Pub., NY.
- 18** Bent, A.C. 1963. Life Histories of North American Wood Warblers, pt 1. Dover Pub., NY.
- 19** Bent, A.C. 1963. Life Histories of North American Wood Warblers, pt 2. Dover Pub., NY.
- 20** Bent, A.C. 1968. Life Histories of North American Cardinals, Grosbeaks, Buntings, Towhees, Finches, Sparrows, and their allies, pt. 1. Dover Pub., NY.
- 21** Bent, A.C. 1968. Life Histories of North American Cardinals, Grosbeaks, Buntings, Towhees, Finches, Sparrows, and their allies, pt. 2. Dover Pub., NY.
- 22** Bent, A.C. 1968. Life Histories of North American Cardinals, Grosbeaks, Buntings, Towhees, Finches, Sparrows, and their allies, pt. 3. Dover Pub., NY.
- 23** Bent, A.C. 1968. Life Histories of North American Wagtails, Shrikes Vireos, and their allies. Dover Pub., NY.
- 24** Bent, A.C. 1963. Life Histories of North American Gulls and Terns. Dover Pub., NY.
- 25** Cahalane, V.H. 1961. Mammals of North America. Macmillan Company, NY.
- 26** Bent, A.C. 1963. Life Histories of North American Marsh Birds. Dover Pub., NY.
- 27** Bent, A.C. 1962. Life Histories of North American Wild Fowl, pt. 1. Dover Pub., NY.
- 28** Bent, A.C. 1962. Life Histories of North American Wild Fowl, pt. 2. Dover Pub., NY.
- 29** Godin A.J. 1977. Wild Mammals of New England. Johns Hopkins University Press, Baltimore, Maryland.

- 30** Bent, A.C. 1962. Life Histories of North American Shore Birds, pt. 1. Dover Pub., NY.
- 31** Bent, A.C. 1962. Life Histories of North American Shore Birds, pt. 2. Dover Pub., NY.
- 32** Bull, J. 1974. Birds of New York State. Doubleday/Natural History Press, Garden City.
- 33** Wright, A.H., and A.A. Wright. 1949. Handbook of Frogs & Toads Comstock Pub. Ass., Ithaca, NY.
- 34** Noble, G.K. 1954. The Biology of the Amphibians, Dover Pub., NY.
- 35** Mattison, C. 1987. Frogs & Toads of the world. Facts On File Pub., NY.
- 36** Bishop, S.C. 1943. Hand Book of Salamanders. Comstock Pub. Ass. Ithaca.
- 37** Dickerson, M.C. 1943. The Frog Book. Dover Pub., NY.
- 38** Leviton, A.E. Reptiles and Amphibians of North America. Doubleday & Company, NY.
- 39** Wright, A.H., and A.A. Wright. 1957. Handbook of Snakes V. 1. Comstock Pub. Ass., Ithaca, NY.
- 40** Wright, A.H., and A.A. Wright. 1957. Handbook of Snakes V. 1. Comstock Pub. Ass., Ithaca, NY.
- 41** Obst, F.J. Turtles, Tortoises, and Terrapins. Saint Martin's Press NY.
- 42** Stone, W. 1965. Bird Studies at Old Cape May V. 1. Dover Pub., NY.
- 43** Stone, W. 1965. Bird Studies at Old Cape May V. 2. Dover Pub., NY.
- 44** Forbush, E.H. 1912. The History of The Game Birds, Wildfowl, and Shore Birds of Massachusetts and Adjacent States. Wright & Potter Printing, Massachusetts.
- 45** Barbour, R.W., and W.H. Davis. 1969. Bats of America. The University Press of Kentucky, Lexington, KY.